


Nachrichten des Entomologischen Vereins Apollo



- 97 LEWANDOWSKI, S., & FISCHER, H.: Zweiter Nachtrag zu den Revisionen der *trifolii*- und *serrula*-Artengruppen der Gattung *Lasiocampa* sowie neue Erkenntnisse zum Typenfundort von *L. decolorata* (Lepidoptera: Lasiocampidae)
- 105 SMETACEK, P.: A new subspecies of *Mycalesis suaveolens* WOOD-MASON & DE NICÉVILLE 1883 from the western Himalaya, India (Lepidoptera, Nymphalidae, Satyrinae)
- 109 SMETACEK, P.: Preimaginal stages of *Graphium (Pazala) eurous caschmirensis* (ROTHSCHILD 1895) (Lepidoptera: Papilionidae) in the Kumaon Himalaya, India
- 113 SMETACEK, P., & KITCHING, I. J.: The hawkmoths of Ladakh, Jammu & Kashmir, India (Lepidoptera: Sphingidae)
- 116 Personalia: Nachträge zur Bibliographie von Werner THOMAS, 8
- 117 NAUMANN, S., NÄSSIG, W. A., & NOGUEIRA G., G.: *Mexicantha oaxacana* gen. et sp. n., a new hemileucine moth from Oaxaca, Mexico (Lepidoptera: Saturniidae)
- 121 ČERNÝ, K.: *Lyclene weidenhofferi* sp. n. and *Barsine delineata* (WALKER, 1854) discovered in Thailand (Lepidoptera: Noctuoidea, Erebidae, Arctiinae, Lithosiini)
- 123 Entomologische Praxis: SCHINTLMEISTER, A.: Der Einsatz von Energiesparlampen für den Lichtfang
- 125 TARMANN, G. M.: Two new species of Chalcosiinae from South East Asia (Lepidoptera: Zygaenidae)
- 130 Buchbesprechung: HOWSE, P., & WOLFE, K. (2011): Giant Silkmoths. Colour, mimicry & camouflage.
- 131 MIELKE, C. G. C., & GREHAN, J. R.: Catalogue of the Latin American Hepialidae with taxonomic remarks (Lepidoptera)
- 159 EFETOV, K. A., GORBUNOV, O. G., & TARMANN, G. M.: Zygaenidae of Mongolia (Lepidoptera)
- 165 BREITHAUPT, R.: Die Raupe von *Ilema costalis* (WALKER, 1855) (Lepidoptera, Lymantriidae)
- 167 Nekrolog: Rudolf E. J. LAMPE, 12. IV. 1933–12. VII. 2011
- 170 Buchbesprechung: WESTENBERGER, A. & FABIAN, J. (2009): Die Tagfalter des Main-Taunus-Kreises und der näheren Umgebung, ihre Raupen und Futterpflanzen.
- 171 Hessenfauna: 28. NÄSSIG, W. A., & ZUB, P. M. T.: Neue Informationen über *Caradrina (Eremodrina) gilva* (DONZEL, 1837) in Hessen (Lepidoptera, Noctuidae)
- 173 ALBRECHT, M.: *Stachys cretica*, a new host plant for *Carcharodus orientalis* REVERDIN, 1913 on the Greek island of Samos (Lepidoptera: Hesperidae)
- 176 Buchbesprechung: RHEINHEIMER, J., & HASSLER, M. (2010): Die Rüsselkäfer Baden-Württembergs.
- 177 BUCHSBAUM, U., DE FREINA, J. J., & CHEN, M.-Y.: *Hyblaea aureola* sp. n. from South Thailand (Lepidoptera, Hyblaeidae) – Studies in Hyblaeidae 2
- 180 Buchbesprechung: WANG, M., & KISHIDA, Y. (2011): Moths of Guangdong Nanling National Nature Reserve
- 181 WEYH, R. E.: Beobachtungen an einer südosthessischen Population von *Chysoesthia drurella* (FABRICIUS, 1775) (Lepidoptera: Gelechiidae, Gelechiinae)
- 185 Entomologische Notiz: SCHURIAN, K. G.: Wasseraufnahme und Wasserabgabe („Mud-puddling“) auch bei *Apatura ilia* ([DENIS & SCHIFFERMÜLLER], 1775) festgestellt (Lepidoptera: Nymphalidae)
- 187 GÖHL, K., BRÜGGEMEIER, F., & THUST, R.: Detection of the forester moth *Adscita capitalis* (STAUDINGER, 1879) (Lepidoptera: Zygaenidae, Procrinae) in Greek Macedonia

Jahrgang 32
Heft 3/4
März 2012

Copyright © 2012 by Entomologischer Verein Apollo e.V., Frankfurt am Main, Germany.

All rights reserved.

No part of this publication may be reproduced in any form, or be used, stored, propagated and distributed using electronic media without written permission of the editor. Authors that receive the pdf version may use it for private exchange with colleagues or for sending on request, only. Online-access in the internet is not allowed except by written permission of the editor.

Alle Rechte vorbehalten.

Kein Teil dieser Publikation darf ohne schriftliche Genehmigung des Herausgebers in irgendeiner Form reproduziert oder unter Verwendung elektronischer Systeme verarbeitet, gespeichert, vervielfältigt oder verbreitet werden. Die Autoren erhalten die pdf-Version nur für den privaten Austausch mit Fachkollegen oder für den Versand auf einzelne Anfragen hin. Sie darf nicht ohne schriftliche Genehmigung des Herausgebers im Internet zugänglich gemacht werden.

ISSN 0723-9912

Catalogue of the Latin American Hepialidae with taxonomic remarks (Lepidoptera)

Carlos G. C. MIELKE and John R. GREHAN

Carlos G. C. MIELKE, Caixa Postal 1206, 84 145-000, Carambeí, PR, Brazil; cmielke1@uol.com.br

Dr. John R. GREHAN, Director of Science and Research, Buffalo Museum of Science, 1020, Humboldt Parkway, Buffalo, NY 14211-1193, USA; jgrehan@sciencebuff.org

Abstract: A catalogue of the Latin American (i.e., from Mexico to the southern end of South America) Hepialidae is presented. Taxonomic comments, type localities, and depository museums are recorded. 125 species in 30 genera are recognized as valid. A checklist is provided at the end.

The following are recognized as **new synonyms**: *Aepytus (Xytrops)* VIETTE, 1951 **syn. n.** of *Cibyra* WALKER, 1856; *Lamelliformia* VIETTE, 1951 **syn. n.** of *Tricladia* C. & R. FELDER, 1874. *Schaefferiana jeanneli* VIETTE, 1950 **syn. n.** of *Aepytus guarani* (PFITZNER, 1914); *Aepytus helga* SCHAUS, 1929 and *Cibyra poltrona* SCHAUS, 1901 **syn. n.** of *Cibyra dorita* SCHAUS, 1901; *Dalaca chiriquensis* PFITZNER, 1914 and *Dalaca muysca* PFITZNER, 1914 **syn. n.** of *Gymelloxos prosopus* (DRUCE, 1901); *Phassus agrionides* WALKER, 1856, *Roseala bourgognei* VIETTE, 1950 and *Thiastyx catharinae* VIETTE, 1951 **syn. n.** of *Roseala tessellatus* (HERRICH-SCHÄFFER, [1854]).

The following are recognized as **new combinations**: *Aepytus guarani* (PFITZNER, 1914), *Cibyra stigmatica* (PFITZNER, 1937), *Gymelloxos prosopus* (DRUCE, 1901), *Pfitzneriana obliquestrigata* (STRAND, 1912), *Philoenia indicata* (STRAND, 1912), *Philoenia thisbe* (DRUCE, 1901), *Pseudodalaca sarta* (SCHAUS, 1894), *Pseudophassus philiponi* (VIETTE, 1950), *Roseala tessellatus* (HERRICH-SCHÄFFER, [1854]), *Tricladia prytanes* (SCHAUS, 1892), *Tricladia sladeni* (HAMPSON, 1903), *Tricladia tupi* (PFITZNER, 1914), *Vietteogorgopis absyrtus* (SCHAUS, 1892), *Vietteogorgopis katharinae* (PFITZNER, 1914), *Yleuxas brunnea* (SCHAUS, 1901).

The following are recognized as **revised combinations**: *Aepytus biedermani* (VIETTE, 1950), *Aepytus exclamans* (HERRICH-SCHÄFFER, [1854]), *Alloaepytus tesselloides* (SCHAUS, 1901), *Cibyra oreas* (SCHAUS, 1892), *Cibyra schausi* (VIETTE, 1952), *Gymelloxos terea* (SCHAUS, 1892), *Gymelloxos trilinearis* (PFITZNER, 1914), *Hampsoniella assa* (DRUCE, 1887), *Hampsoniella equatorialis* (VIETTE, 1950), *Hepialyxodes rileyi* VIETTE, 1951, *Philoenia brasiliensis* VIETTE, 1952, *Philoenia fasslii* (PFITZNER, 1914), *Philoenia guyanensis* (VIETTE, 1951), *Philoenia lagopus* (MÖSCHLER, 1877), *Philoenia saguanmachica* (PFITZNER, 1914), *Pseudodalaca gugelmanni* (VIETTE, 1950), *Pseudodalaca mexicanensis* VIETTE, 1953, *Pseudophassus mahagoniatus* PFITZNER, 1914, *Pseudophilaenia omagua* (PFITZNER, 1937), *Schaefferiana epigramma* (HERRICH-SCHÄFFER, [1854]), *Schaefferiana simplex* VIETTE, 1956, *Tricladia umbrifera* C. & R. FELDER, 1874, *Yleuxas bradleyi* VIETTE, 1951. The following are recognized as **revised status to genus rank**: *Aepytus* HERRICH-SCHÄFFER, [1856], *Alloaepytus* VIETTE, 1951, *Gymelloxos* VIETTE, 1952, *Hampsoniella* VIETTE, 1950, *Hepialyxodes* VIETTE, 1951, *Philoenia* KIRBY, 1892, *Pseudodalaca* VIETTE, 1951, *Pseudophassus* PFITZNER, 1914, *Pseudophilaenia* VIETTE, 1951, *Schaefferiana* VIETTE, 1950, *Tricladia* C. & R. FELDER, 1874, *Yleuxas* VIETTE, 1951, *Tricladia tupi* (PFITZNER, 1914), *Yleuxas bradleyi* VIETTE, 1951.

The following is transferred to another family: *Acrolophus tapuja* (PFITZNER, 1914), **comb. n.** (to Tineidae).

Keywords: taxonomy, Tineidae.

Systematisches Verzeichnis der lateinamerikanischen Hepialidae mit taxonomischen Anmerkungen (Lepidoptera)

Zusammenfassung: Es wird ein systematisches Verzeichnis der lateinamerikanischen (von Mexiko bis zur Südspitze Südamerikas) Hepialidae (Lepidoptera) präsentiert. Taxonomische Anmerkungen, Typenfundorte und Typenhinterlegungsstätten werden aufgeführt. 125 Arten in 30 Gattungen werden als valide anerkannt (siehe Checkliste am Ende).

Die Auflistungen der neuen Synonyme, neuen und revidierten Kombinationen sowie revidierten Statuszuweisungen siehe im Abstract. Ein Arttaxon wurde in eine andere Familie transferiert (siehe Abstract).

Introduction

Latin American (= from Mexico to the southern end of South America) Hepialidae have been poorly understood and studied since the first species was described by GEYER ([1838]). Despite of this neglect the Neotropical Hepialidae fauna can be compared to other regions well treated like Australia considering the number of species (NIELSEN et al. 2000). These authors summarized 616 species in the world, being 134, in 20 genera, for the Neotropical region.

Early documentation of the Latin American Hepialidae was sporadic, beginning with WALKER's (1856) list of specimens in the British Museum (London) and KIRBY's (1882) global catalogue titled "A synonymic catalogue of Lepidoptera Heterocera (moths)". A further three decades were to pass before publication of the chapter on Hepialidae within the "Lepidopterorum Catalogus" by WAGNER & PFITZNER (1911), and in the grandly titled book series "Die Gross-Schmetterlinge der Erde", edited by SEITZ and written by PFITZNER (1937–38), many species were figured for the first time. Over much of the following six decades further taxonomic developments were limited to brief descriptions of new species and genera, particularly by Pierre VIETTE who described several new species and genera in a series of articles from 1949 to 1961. But there were no further catalogues until the taxonomic treatment of Southern South American Hepialidae by NIELSEN & ROBINSON (1983) that included a list for Latin American species, and a similar catalogue produced in the following year by ROBINSON & NIELSEN (1984). These lists were then updated in the most recent global inventory by NIELSEN et al. (2000).

In the present catalogue, references on systematic, geographical distribution, ethology, ecology, biology, and host-plants are provided for each genus, species and sub-

species with the exception of a few articles on pasture and/or agricultural damage that could not be obtained.

Whenever a taxonomic change is made, an explanation is presented at the end of the taxon, with the exception of VIETTE's genera that were designated subgenera by NIELSEN et al. (2000) because there was no justification for the subgeneric change, and in our examination of external features we did not find any features that would justify their placement within *Cibyra*, and given the considerable ♂ genitalic variation among VIETTE's genera, their generic status remains warranted at this time.

NIELSEN & ROBINSON (1983) were the first authors to list all described species for Latin America, after several species were described by VIETTE. Several species originally described in *Dalaca* (a Chilean genus) were categorized by these authors as "*Dalaca*" *sensu lato* to highlight their identity not conforming to *Dalaca sensu stricto*. After examining accessible holotypes we could locate some species within described genera, but the remaining 7 species are presented in this catalogue as *incertae sedis* pending future studies to clarify their taxonomic status. In addition, some species previously placed in recognized genera are also relocated to this section because they do not conform to the type-species or lack sufficient information to recognize their generic association.

Acknowledgments

We are indebted to Dra. Mirna M. CASAGRANDE (Curitiba, Paraná) and Dr. Olaf H. H. MIELKE (Curitiba, Paraná) for critical review of the manuscript, information, bibliography and enthusiasm. Dr. Joël MINET (MNHN, Paris, France), Dr. Wolfgang A. NÄSSIG (SMFL, Frankfurt am Main, Germany), Dr. Sabine GAAL (NHWM, Vienna, Austria), Dr. Wolfram MEY (ZMHB, Berlin, Germany), Ulf BUCHSBAUM and Dr. Axel HAUSMANN (ZSBS, Munich, Germany), Dr. Kevin TUCK and Dra. Blanca HUERTAS (BMNH, London, England) for accessing their institutions and making the examinations possible. Dra. Jane COSTA (Instituto Oswaldo Cruz, Rio de Janeiro, Rio de Janeiro) and Dr. Marcelo DUARTE (Museu de Zoologia de São Paulo, São Paulo) for accessing their institutions and loan of specimens. Dr. Rodolphe ROUGERIE (Rouen, France) and Dr. Thibaud DECÄENS (Rouen, France) for helping by different ways, specially by accessing mtDNA barcode sequences. Abílio RANK†, Ivo RANK and Osiel RANK (São Bento do Sul, Santa Catarina), Alfred MOSER (São Leopoldo, Rio Grande do Sul), Antônio PESSOA (Sobral, Ceará), Daniel HERBIN (Toulouse, France), Eduardo A. PEREIRA (Conceição dos Ouros, Minas Gerais), Eurides FURTADO (Diamantino, Mato Grosso), Guilherme P. DE ALMEIDA NETO (Conceição dos Ouros, Minas Gerais), Herbert MIERS† (Joinville, Santa Catarina), Jean HAXAIRE (Laplume, France), José MONZÓN (Guatemala), Nirton TANGERINI (Rio de Janeiro), Ricardo SIEWERT (Pelotas, Rio Grande do Sul), Thierry PORION (Jaujac, France), Dr. Vitor O. BECKER (Camacã, Bahia) for supplying field information and specimens. Luis DEVOTTO and Michel

DOTTAX for bibliography. Special thanks to Elyana G. JOERKE (Ponta Grossa, Paraná) and Ernst BROCKMANN (Lich, Germany) for helping in various ways, and Pierre JAUFFRET† (Santo Antônio do Tauá, Pará) for translating many of VIETTE's articles.

Systematic part

Acronyms

Collections housing primary types are mentioned by the following acronyms:

- BMNH The Natural History Museum, London, England (visited by CGCM).
- MACN Museo Argentino de Ciencias Naturales Bernardino Rivadavia, Buenos Aires, Argentina.
- MNHC Museo Nacional de Historia Natural, Santiago, Chile.
- MNHN Muséum National d'Historie Naturelle, Paris, France (visited by CGCM).
- NHMW Naturhistorisches Museum, Vienna, Austria (visited by CGCM).
- SMFL Senckenberg-Museum, Lepidoptera collection, Frankfurt am Main, Germany (visited by CGCM).
- SMFT Type catalogue number of Senckenberg-Museum, Lepidoptera collection, Frankfurt am Main, Germany.
- NRSS Naturhistoriska Riksmuseet, Stockholm, Sweden.
- RMNH Naturalis, formerly Rijksmuseum van Natuurlijke Historie, Leiden, Holland.
- USNM United States National Museum of Natural History, Washington DC, USA (visited by CGCM).
- ZMHB Museum für Naturkunde der Humboldt-Universität, Berlin, Germany (visited by CGCM).
- ZSBS Zoologische Sammlungen des bayerischen Staates, Munich, Germany (visited by CGCM).
- ZMUC Zoological Museum, Copenhagen, Denmark.

Further abbreviations:

- ◇ Indicating the same author[s] as the one[s] just before (name[s] not repeated).
- gen. Genitalia figure.
- GP Genitalia preparation/dissection/slide number.

Aepytus HERRICH-SCHÄFFER, [1856] **stat. rev.**

Type-species: *Epialus* [sic] *exclamans* HERRICH-SCHÄFFER, [1854], by subsequent designation by KIRBY (1892: 887).

Aepytus HERRICH-SCHÄFFER ([1856]: 5); included species: *exclamans* HERRICH-SCHÄFFER, [1854], *costalis* HERRICH-SCHÄFFER, [1854], *nanus* HERRICH-SCHÄFFER, [1854], *catocalus* HERRICH-SCHÄFFER, [1854].

HERRICH-SCHÄFFER ([1858]: 56), syn.: *Dalaca* WALKER, 1856 – NEAVE (1939: 75). – VIETTE (1950a: 73); ◇ (1951b: 116); ◇ (1951d: 74). – PAULT (1953: 143), syn.: *Alloaepytus* VIETTE, 1951, *Hampsoniella* VIETTE, 1949, *Hepialyxodes* VIETTE, 1951, *Parana* VIETTE, 1949, *Philoenia* KIRBY, 1892, *Pseudodalaca* VIETTE, 1949, *Pseudophilaenia* VIETTE, 1950, *Yleuxas* VIETTE, 1951; ◇ (1957: 51), syn.: *Gymelloxos* VIETTE, 1952, *Pfitzneriana* VIETTE, 1952, *Aplatissa* VIETTE, 1953. – NIELSEN & ROBINSON (1983: 19). – WINDER & HARLEY (1983: 355). – ROBINSON & NIELSEN (1984: 17). – GREHAN (1989: 805). – NYE & FLETCHER (1991: 8). – HILJE et al. (1992b: 152–154; fig. 54 [larva]). – HILJE et al. (1992a: 152). – KRISTENSEN (1998: 62). – NAIR (2001: 21). – GREHAN & RAWLINS (2003: 734); as synonym

of *Cibyra* WALKER, 1856. — MACÍAS et al. (2005: 100). — ARGUEDAS (2004: 6; fig. 7a, b [larva]); ◇ (2006: 7); ◇ (2007: 9, 28, 57).

Aepytus (*Aepytus*): VIETTE (1950a: 74). — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Aepytus*) WALKER, 1856: NIELSEN et al. (2000: 842). — SIMONSEN (2002: 65). — GREHAN (2010: 49).

Remarks. Based on the morphology of the type-species, the genus *Aepytus* HERRICH-SCHÄFFER, [1856] is revalidated. This genus shows “oxycanine” (see DUMBLETON 1966: 922) wing venation.

Aepytus biedermanni (VIETTE, 1950), **comb. rev.**

Schaefferiana biedermanni VIETTE (1950c: 60; fig. 10 valva): [holo-]type ♂, Brazil meridional, Sud de l'Etat de Minas Geraes [RECTE Gerais], [Catas Altas], Caraça, 2^o semestre 1884, P. GERMAIN [leg.]; coll. R. BIEDERMANN, ex coll. C. OBERTHÜR; GP VIETTE 1365; MNHN.

Aepytus biedermanni: VIETTE (1951c: 96).

Aepytus (*Aepytus*) *biedermanni*: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Aepytus*) *biedermanni*: NIELSEN et al. (2000: 842).

Aepytus exclamans (HERRICH-SCHÄFFER, [1854]), **comb. rev.**

Epialus [SIC] *exclamans* HERRICH-SCHÄFFER ([1854] (BOISDUVAL, in litt.): cover; pl. [31], fig. 145 [♂ dorsal]); Brazil; [ex coll. BOISDUVAL]; [GP P. VIETTE no. 2326]; [MNHN]; ◇ ([1858]: 79).

Dalaca exclamans: WALKER ([v.] 1856: 1561). — GERSTAECKER (1857: 425). — WAGNER & PFITZNER (1911: 13). — PFITZNER (1937: 1293; pl. 185a [♂ dorsal]).

Aepytus exclamans: HERRICH-SCHÄFFER ([ix. 1856]: 5); ◇ ([1858]: 56). — KIRBY (1892: 887). — VIETTE (1951c: 95).

Aepytus (*Aepytus*) *exclamans*: VIETTE (1950a: 75; fig. 1 ♂ gen. [error, *Cibyra dormita* SCHAUS, 1901]). — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Aepytus*) *exclamans*: NIELSEN et al. (2000: 842).

Aepytus guarani (PFITZNER, 1914), **comb. n.**

Dalaca guarani PFITZNER (1914: 105): Southern Brazil, St. Katharina (RECTE Santa Catarina); coll. PFITZNER; [SMFL]; ◇ (1937: 1296; pl. 100b, [♀] dorsal). — VIETTE (1952a: 144); fig. 2 (♀ gen.). — SCHRÖDER (1967: 339; “holotype” [RECTE lectotype] [♀], SMFT 90). — NIELSEN & ROBINSON (1983: 20, 44). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

= *Schaefferiana jeanneli* VIETTE (1950c: 59; fig. 8, ♂ gen.): [holo-]type ♂, Brazil, Paraná, Curitiba [RECTE Curitiba], ix. 1911, P. LOMBARD [leg.]; GP P. VIETTE 951; MNHN; **syn. n.**

Aepytus jeanneli: VIETTE (1951c: 96); ◇ (1951e: 1275).

Aepytus (*Aepytus*) *jeanneli*: NIELSEN & ROBINSON (1983: 19, 44). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Aepytus*) *jeanneli*: NIELSEN et al. (2000: 842). — SIMONSEN (2002: 65).

Dalaca (*Aepytus*) *guarani* [SIC]: GREHAN (2010: 51; fig. a, appendix).

Remarks. *D. guarani* PFITZNER, 1914 was described based on the ♀ and *S. jeanneli* VIETTE, 1950 on the ♂ of the same species. Matching was done through examination of the types, morphology, and also through mtDNA barcode sequences.

Alloaepytus VIETTE, 1951, **stat. rev.**

Type-species: *Dalaca tesselloides* SCHAUS, 1901 by original designation.

Aepytus (*Alloaepytus*) VIETTE (1951a: 2); included species: *tesselloides* (SCHAUS, 1901), *coscinophora* (PFITZNER, 1914).

EDWARDS & HOPWOOD (1966: 9). — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Alloaepytus: VIETTE (1956: 378). — NYE & FLETCHER (1991: 12).

Cibyra (*Alloaepytus*): NIELSEN et al. (2000: 842). — GREHAN (2010: 49).

Remarks. PAULT (1953: 145): as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. Based on the genitalia morphology of the type-species, *Alloaepytus* VIETTE, 1951 is revalidated at full genus rank.

Alloaepytus tesselloides (SCHAUS, 1901), **comb. rev.**

Dalaca tesselloides SCHAUS (1901: 76): [♀], Paraguay; [GP VIETTE no. 91519, type no. USNM 18610]; USNM. — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1296). — ZUKOWSKI (1954: 93), **syn.:** *D. stigmatica* PFITZNER.

= *Dalaca coscinophora* PFITZNER (1914: 105): [Brazil], Mato-Grosso [RECTE Mato Grosso do Sul], Nivac [RECTE Nioaque]; coll. PFITZNER; [SMFL]; ◇ (1937: 1294; pl. 100b, [♂] dorsal). — SCHRÖDER (1967: 339): lectotype [♂], SMFT 83.

Aepytus (*Alloaepytus*) *coscinophora*: VIETTE (1951a: 4). — NIELSEN & ROBINSON (1983: 19): as synonym of *Aepytus* (*Alloaepytus*) *tesselloides* (SCHAUS, 1901). — ROBINSON & NIELSEN (1984: 17): as synonym of *Aepytus* (*Alloaepytus*) *tesselloides* (SCHAUS, 1901).

Aepytus (*Alloaepytus*) *tesselloides*: VIETTE (1951a: 4; fig. 2, ♂ gen.). — NIELSEN & ROBINSON (1983: 19), **syn.:** *Aepytus* (*Alloaepytus*) *coscinophora* (PFITZNER, 1914). — ROBINSON & NIELSEN (1984: 17), **syn.:** *Aepytus* (*Alloaepytus*) *coscinophora* (PFITZNER, 1914).

Alloaepytus coscinophora: VIETTE (1956: 378).

Alloaepytus tesselloides: VIETTE (1956: 379).

Cibyra (*Alloaepytus*) *coscinophora*: NIELSEN et al. (2000: 842) as synonym of *Cibyra* (*Alloaepytus*) *tesselloides* (SCHAUS, 1901).

Cibyra (*Alloaepytus*) *tesselloides*: NIELSEN et al. (2000: 842): **syn.:** *Cibyra* (*Alloaepytus*) *coscinophora* (PFITZNER, 1914). — GREHAN (2010: 51; fig. b, appendix).

Andeabatis NIELSEN & ROBINSON, 1983

Type-species: *Xyleutes chilensis* URETA, 1951, by original designation; monotypic.

NIELSEN & ROBINSON (1983: 108). — ROBINSON & NIELSEN (1984: 16). — NYE & FLETCHER (1991: 19). — EDWARDS et al. (1996: 34). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN (2010: 49).

Andeabatis chilensis (URETA, 1951)

Xyleutes chilensis URETA (1951: 75; 2 figs.: ♀ [RECTE ♂] dorsal, ventral; holotype ♀ [RECTE ♂], Chile, [Malleco], Curacautín, Termas de Río Blanco, ii. 1946 (RECTE 1944; see NIELSEN & ROBINSON 1983: 110), A. WAGENKNECHT leg.; MNHC. — CAMOUSSEIGHT (1980: 31).

Phassus chilensis: URETA (1956: 282). — MALLET (1984: 77).

Andeabatis chilensis: NIELSEN & ROBINSON (1983: 110; figs. 71 (prelabium, labial palpus), 72 (venation), 73 (forewing scale), 212 (♂ dorsal), 252 (♂ antennae flagellum), 277 (♂ foretibia), 333 (♂ gen.), 363 (♀ gen.), 415 (flight period), 431 (distribution)): type no. 705. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — PASTRANA (2004: 5).

Aplatissa VIETTE, 1953

Type-species: *Aplatissa strangoides* VIETTE, 1953, by original designation; monotypic.

VIETTE (1953b: 81) — PACLT (1957: 51): as synonym of *Aepytus* HERRICH-SCHÄFFER, [1858]. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NYE & FLETCHER (1991: 26). — NIELSEN et al. (2000: 841). — GREHAN (2010: 49).

Aplatissa michaelis (PFITZNER, 1914)

Dalaca michaelis PFITZNER (1914: 105): a pair, Peru, Chanchamayo, 1000 m; single specimen, high Amazonas, Otto MICHAEL leg.; coll. PFITZNER; [SMFL]. — SCHRÖDER (1967: 340): lectotype ♂ [♀?], SMFT 87.

Dalaca michaeli [sic]: PFITZNER (1937: 1294; pl. 99 d, [♂, ♀] dorsal). — ZUKOWSKI (1954: 93).

Aplatissa michaelis: NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

Aplatissa strangoides VIETTE, 1953

Aplatissa strangoides VIETTE (1953b: 81; fig. 1 (♂ gen.)): holotype ♂, Brazil, [Amazonas], Fonte Boa, vi. 1906, S. M. KLAGES leg.; GP P. VIETTE no. 2414; BMNH. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

Blanchardinella NIELSEN, ROBINSON & WAGNER, 2000

Type-species: *Hepialus venosus* BLANCHARD, 1852, by original designation by VIETTE (1950e: 145) (replacement name).

Blanchardina VIETTE (1950e: 145), monotypic; **praecoc.**: *Blanchardina* LABBÉ, 1899 [Protozoa]. — PACLT (1953: 145): as synonym of *Dalaca* WALKER, 1856. — EDWARDS & HOPWOOD (1966: 37). — NIELSEN & ROBINSON (1983: 17, 88). — ROBINSON & NIELSEN (1984: 16). — NYE & FLETCHER (1991: 45). — NIELSEN et al. (2000: 840).

Blanchardinella NIELSEN et al. (2000: 823, 840): replacement name for *Blanchardina* VIETTE, 1950. — SIMONSEN (2002: 65). — GREHAN (2010: 49).

Blanchardinella venosus (BLANCHARD, 1852)

Hepialus venosus BLANCHARD (1852: 70; pl. 4, fig. 6 [♂ dorsal]); Chile, Coquimbo; [MNHN]. — WALKER (1856: 1557).

Dalaca venosa [sic]: BUTLER (1882: 25), syn.: *Dalaca nigricornis* WALKER, 1856. — AURIVILLIUS (1884: 524), syn.: *Dalaca nigricornis* WALKER, 1856. — KIRBY (1892: 886), syn.: *Dalaca nigricornis* WALKER, 1856. — WAGNER & PFITZNER (1911: 14), syn.: *Dalaca nigricornis* WALKER, 1856. — PFITZNER (1937: 1295), syn.: *Dalaca nigricornis* WALKER, 1856. — PACLT (1949: 149), syn.: *Dalaca parviguttata* (BRYK, 1944); ♀ (1953: 146), syn.: *Dalaca parviguttata* (BRYK).

Blanchardina venosus: VIETTE (1950e: 145; fig. 4, [syn-]type ♂ gen., GP VIETTE no. 939). — NIELSEN & ROBINSON (1983: 17, 90; figs. 54 (prelabium and labial palpus), 55 (venation), 187–188 (♂ dorsal), 189–191 (♀ dorsal), 244 (♂ antennae flagellum), 268 (♂ foretibia), 269 (♀ foretibia), 323–324 (♂ gen.), 355a, b (♀ gen.), 380 (bursa copulatrix), 407 (flight period), 428 (distribution)). — ROBINSON & NIELSEN (1984: 16).

Blanchardinella venosus: NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN (2010: 60; fig. M, appendix).

Remarks. *Dalaca (Triodia) venosa* [sic] BERG, 1882 is a misidentification; see *Dalaca pallens* (BLANCHARD, 1852), NIELSEN & ROBINSON (1983: 90).

Calada NIELSEN & ROBINSON, 1983

Type-species: *Calada fuegensis* NIELSEN & ROBINSON, 1983, by original designation.

Calada NIELSEN & ROBINSON (1983: 17, 91); included species: *fuegensis* NIELSEN & ROBINSON, 1983, *migueli* NIELSEN & ROBINSON, 1983.

ROBINSON & NIELSEN (1984: 16). — NYE & FLETCHER (1991: 51). — EDWARDS et al. (1996: 107). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN (2010: 49).

Calada fuegensis NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 93; figs. 56 (prelabium and labial palpus), 57 (venation), 192–195 (♂ dorsal), 196 (♀ dorsal), 245 (♂ antennae flagellum), 270 (♂ foretibia), 271 (♀ foretibia), 325 (holotype ♂ gen.), 356 (♀ gen.), 381 (bursa copulatrix), 408 (flight period), 429 (distribution)); holotype ♂, Argentina, Tierra del Fuego, Isla Grande, W of Ushuaia, Lapataia, 20 m, 2. ii. 1979, Mision Cientifica Danesa leg., sta. 34; GP ESN 2740; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — PASTRANA (2004: 5).

Calada migueli NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 94; figs. 197 (holotype ♂ dorsal), 272 (♂ foretibia), 326 (♂ gen.), 409 (flight period), 429 (distribution)); holotype ♂, Argentina, Rio Negro, Lago Nahuel Huapi, Porto Blest, 770 m, 2. iii. 1979, Mision Cientifica Danesa leg., sta. 8; GP ESN 2548; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — PASTRANA (2004: 5).

Callipielus BUTLER, 1882

Type-species: *Callipielus arenosus* BUTLER, 1882 by monotypy.

BUTLER (1882: 23). — KIRBY (1892: 893). — WAGNER & PFITZNER (1911: 22). — PFITZNER (1938: 1298). — NEAVE (1939: 536). — BRYK (1944: 26). — PACLT (1944: 143). — VIETTE (1950c: 52); ♀ (1951d: 74). — PACLT (1953: 143). — ROBINSON (1977: 110; figs. 1 (♂ gen.), 2 (aedeagus), 3 (caudal sclerites), 4 (venation)), syn.: *Stachyocera* URETA, 1957. — NIELSEN & ROBINSON (1983: 17, 73), syn.: *Stachyocera* URETA, 1957. — MALLET (1984: 77). — ROBINSON & NIELSEN (1984: 16), syn.: *Stachyocera* URETA, 1957. — NIELSEN & SCOBLE (1986: 43). — NYE & FLETCHER (1991: 52). — KRISTENSEN (1998: 62). — NIELSEN et al. (2000: 839), syn.: *Stachyocera* URETA, 1957. — SIMONSEN (2002: 65). — GREHAN (2010: 44).

= *Stachyocera* URETA (1957: 159); type-species: *Stachyocera izquierdoi* URETA, 1957, by original designation. — EDWARDS & VEVEES (1975: 325). — ROBINSON (1977: 110); as synonym of *Callipielus* BUTLER, 1882. — NIELSEN & ROBINSON (1983: 17, 73); as synonym of *Callipielus* BUTLER, 1882. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus* BUTLER, 1882. — NYE & FLETCHER (1991: 285). — NIELSEN et al. (2000: 839); as synonym of *Callipielus* BUTLER, 1882.

Callipielus [sic]: DUMBLETON (1966: 924).

Callipielus arenosus BUTLER, 1882

Callipielus arenosus BUTLER (1882: 24; pl. 1, fig. 6 venation); [Chile], Valdivia, coll. REED; [BMNH]. — AURIVILLIUS (1884: 524). — KIRBY (1892: 893). — WAGNER & PFITZNER (1911: 22). — PFITZNER (1938: 1298). — VIETTE (1950c: 52; figs. 1–2 ♂ gen.). — URETA (1957: pl. 1, fig. 9 ([♂] dorsal)). — ROBINSON (1977: 112; pl. 1, fig. 1 (♂ dorsal), pl. 2, fig. 10 (♂ gen., holotype of *Callipielus leukogramma* BRYK, 1944), pl. 3, fig. 18 (caudal sclerites, holotype of *Callipielus leukogramma* BRYK, 1944), pl. 5, figs. 26 (aedeagus, holotype of *Callipielus leukogramma* BRYK, 1944), 28 (♀ gen.), pl. 6, fig. 30 (antennal segments, lectotype)), lectotype ♂ designated; GP BMNH Hepial. 12348, syn.: *Callipielus leukogramma* BRYK, 1944. — NIELSEN & ROBINSON (1983: 17, 76; figs. 159–161 and 163–164 (♂ dorsal), 162 and 165 (♀ dorsal), 232 (♂ antennae flagellum), 233 (♀ antennae flagellum), 262 (♂ foretibia), 263 (♀ foretibia), 303–305 (♂ gen.), 306 (holotype ♂ gen. of *Callipielus leukogramma* BRYK, 1944), 349–350 (♀ gen.), 374 (bursa

copulatrix), 397 (flight period), 424 (distribution)), syn.: *Callipielus antarcticus* (STAUDINGER, 1899) (NEC WALLENGREN, 1860), *Callipielus staudingeri* (WAGNER, 1911), *Callipielus leukogramma* BRYK, 1944, *Callipielus chiliensis* VIETTE, 1950, *Callipielus antarctica* [sic] (STAUDINGER, 1899) (NEC WALLENGREN, 1860). — ROBINSON & NIELSEN (1984: 16), syn.: *Callipielus antarcticus* (STAUDINGER, 1899) (NEC WALLENGREN, 1860), *Callipielus staudingeri* (WAGNER, 1911), *Callipielus leukogramma* BRYK, 1944, *Callipielus chiliensis* VIETTE, 1950. — NIELSEN et al. (2000: 839), syn.: *Callipielus antarcticus* (STAUDINGER, 1899) (NEC WALLENGREN, 1860), *Callipielus staudingeri* (WAGNER, 1911), *Callipielus leukogramma* BRYK, 1944, *Callipielus chiliensis* VIETTE, 1950. — SIMONSEN (2002: 66; figs. 6–10, scales). — DAPOTO et al. (2003: 100). — PASTRANA (2004: 5). — GREHAN (2010: 61; fig. P, appendix).

= *Hepialus antarcticus* STAUDINGER (1899: 42); one ♀ [holotype], [Argentina], Ostküste Feuerlands [East Tierra del Fuego], [Peninsula El] Paramo, North of Sebastians-Bay, 2. i. 1896, OHLIN leg.; [NRSS], praecoc.: *Hepialus antarcticus* WALLENGREN, 1860 [Hepialidae]. — PAGENSTECHER (1902: 399). — WAGNER & PFITZNER (1911: 10). — ENDERLEIN (1912: 90).

= *Hepialus staudingeri* WAGNER (1911: 10): replacement name for *Hepialus antarcticus* STAUDINGER.

Hepialus (*Hepialus*) *staudingeri*: PFITZNER (1937: 1290).

= *Callipielus leukogramma* BRYK (1944: 26; pl. 2, figs. 15 (♂ dorsal), 16 (♀ dorsal)); holotype ♂, [Argentina, Rio Negro, San Carlos de Bariloche], Nahuel Huapi, Peninsula Llau Llau, N of Puerto Nuevo, Patagonia, 770–780 m, x. 1933–iii. 1934, LJUNGNER leg.; NRSS. — VIETTE (1950c: 52). — ROBINSON (1977: 112); holotype GP GSR S-02; as synonym of *Callipielus arenosus* BUTLER, 1882. — NIELSEN & ROBINSON (1983: 17, 76; fig. 306, holotype gen.); as synonym of *Callipielus arenosus* BUTLER, 1882. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus arenosus* BUTLER, 1882. — NIELSEN et al. (2000: 840); as synonym of *Callipielus arenosus* BUTLER, 1882.

= *Callipielus chiliensis* VIETTE (1950f: 74; figs. 16 ♂ dorsal, 18–19 ♂ gen.); [holo-]type ♂, Chile, Valdivia, i. 1938, ANDREAS & DE GRAAG; GP P. VIETTE no. 2187; RMNH. — ROBINSON (1977: 113; pl. 1, fig. 2 (♂ dorsal), pl. 2, fig. 11 (♂ gen.), pl. 3, fig. 19 (caudal sclerites)). — NIELSEN & ROBINSON (1983: 17, 76); as synonym of *Callipielus arenosus* BUTLER, 1882. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus arenosus* BUTLER, 1882. — NIELSEN et al. (2000: 840); as synonym of *Callipielus arenosus* BUTLER, 1882.

Callipielus antarcticus: NIELSEN & ROBINSON (1983: 17, 76); as synonym of *Callipielus arenosus* BUTLER, 1882. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus arenosus* BUTLER, 1882. — NIELSEN et al. (2000: 839); as synonym of *Callipielus arenosus* BUTLER, 1882.

Callipielus antarctica [sic]: NIELSEN & ROBINSON (1983: 17); as synonym of *Callipielus arenosus* BUTLER, 1882.

Callipielus staudingeri: NIELSEN & ROBINSON (1983: 17, 76); as synonym of *Callipielus arenosus* BUTLER, 1882. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus arenosus* BUTLER, 1882. — NIELSEN et al. (2000: 840); as synonym of *Callipielus arenosus* BUTLER, 1882.

Remarks. *Callipielus arenosus* BUTLER, 1882 (*in* PFITZNER 1938) is a misidentification (only figure); see *Callipielus digitata* ROBINSON, 1977, ROBINSON (1977: 115).

Callipielus argentata URETA, 1957

URETA (1957: 162; fig. 3, ♂ gen.); holotype ♂, Chile, [Talca], Laguna del Maule, 1800 m, 31. iii. 1957, G. MARCHANT leg.; coll. M.N. no. 6203; MNHC. — ROBINSON (1977: 118; pl. 1, fig.

8 (♂ dorsal), pl. 3, fig. 17 (♂ gen.), pl. 4, fig. 25 (8th caudal sclerites), pl. 6, fig. 35 (antennal segments)). — CAMOUSSEIGHT (1980: 31). — NIELSEN & ROBINSON (1983: 17, 84; figs. 177 (♂ dorsal), 240 (♂ antennal flagellum), 318 (♂ gen.), 403 (flight period), 427 (distribution)); type no. 1050. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

Callipielus digitata ROBINSON, 1977

ROBINSON (1977: 114; pl. 1, fig. 3 (holotype dorsal), pl. 2, fig. 12 (holotype gen.), pl. 4, fig. 20 (holotype caudal sclerites), pl. 6, figs. 31 (holotype antennal segments), 38 (holotype labium, labial palpus); holotype ♂, Chile; GP GSR H-01; ZMHB. — NIELSEN & ROBINSON (1983: 17, 79; figs. 166–167 (♂ dorsal), 168 (♀ dorsal), 234 (♂ antennae flagellum), 264 (♂ foretibia), 265 (♀ foretibia), 307–309 (♂ gen.), 310 (holotype ♂ gen. of *C. castilloi* ROBINSON, 1977), 351 (♀ gen.), 375 (bursa copulatrix), 398 (flight period), 425 (distribution)), syn.: *Callipielus brunnescens* ROBINSON, 1977, *Callipielus castilloi* ROBINSON, 1977. — ROBINSON & NIELSEN (1984: 16), syn.: *Callipielus brunnescens* ROBINSON, 1977, *Callipielus castilloi* ROBINSON, 1977. — NIELSEN et al. (2000: 840), syn.: *Callipielus brunnescens* ROBINSON, 1977, *Callipielus castilloi* ROBINSON, 1977.

Callipielus arenosus [misidentification]: PFITZNER (1938; pl. 185e, [♂] dorsal); see ROBINSON (1977: 115).

= *Callipielus brunnescens* ROBINSON (1977: 115; pl. 1, fig. 4 (holotype dorsal), pl. 2, fig. 13 (holotype gen.), pl. 4, fig. 21 (holotype caudal sclerites), pl. 6, fig. 32 (holotype antennal segments)); holotype ♂, Chile, LOSSBERG leg.; GP GSR H-02; ZMHB. — NIELSEN & ROBINSON (1983: 17, 79); as synonym of *Callipielus digitata* ROBINSON, 1977. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus digitata* ROBINSON, 1977. — NIELSEN et al. (2000: 840); as synonym of *Callipielus digitata* ROBINSON, 1977.

= *Callipielus castilloi* ROBINSON (1977: 116; pl. 1, fig. 5 (holotype dorsal), pl. 3, fig. 14 (holotype gen.), pl. 4, fig. 22 (holotype caudal sclerites)); holotype ♂, Chile, Cautin, Temuco, Carillanca Experimental Station, 3. vii. 1975, SALAS leg.; gen. slide no. 13147; BMNH. — NIELSEN & ROBINSON (1983: 17, 79; fig. 310 (holotype ♂ gen.)); as synonym of *Callipielus digitata* ROBINSON, 1977. — ROBINSON & NIELSEN (1984: 16); as synonym of *Callipielus digitata* ROBINSON, 1977. — NIELSEN et al. (2000: 840); as synonym of *Callipielus digitata* ROBINSON, 1977.

Callipielus digitatus [sic]: PASTRANA (2004: 5).

Callipielus fumosa NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 83; figs. 176 (holotype dorsal), 239 (♂ antennae flagellum), 317 (holotype gen.), 402 (flight period), 426 (distribution)); holotype ♂, Chile, Nuble, SW side of Vulcan Chillan, Shangri-la, 1600 m, 19.–21. i. 1979, D. & M. DAVIS, AKERBERGS leg.; GP GSR 1044, type no. 100597; USNM. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

Callipielus gentili NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 82; figs. 174–175 (♂ dorsal), 238 (♂ antennae flagellum), 315 (♂ gen.), 316 (holotype gen.), 401 (flight period), 426 (distribution)); holotype ♂, Argentina, Neuquen, Paso Carrirrine, 1000 m, 15. ii. 1966, GENTILI leg.; GP GSR 1033; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — PASTRANA (2004: 5).

Callipielus izquierdoi (URETA, 1957)

Stachyocera izquierdoi URETA (1957: 159; figs. 1 (venation), 2 (♂ gen.), pl. 1, fig. 10 (♂ dorsal)); holotype ♂, Chile, [Arauco], Caramávida, 1000 m, Cordillera de Nahuelbuta, 5. ii.

1953, Luis PeÑA leg.; coll. M.N. no. 6158; MNHC. — CAMOUSEIGHT (1980: 31).

Callipielus izquierdoi: ROBINSON (1977: 117; pl. 1, fig. 7 (♂ dorsal), pl. 3, fig. 16 (♂ gen.), pl. 4, fig. 24 (8th caudal sclerites), pl. 5, fig. 29 (♀ gen.), pl. 6, fig. 34 (antennal segments)). — NIELSEN & ROBINSON (1983: 17, 85; figs. 179–180 (♂ dorsal), 181 (♀ dorsal), 242 (♂ antennae flagellum), 319–320 (♂ gen.), 353b (♀ gen.), 378 (bursa copulatrix), 405 (flight period), 427 (distribution)); type no. 1048. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

Callipielus kraehmeri NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 85; figs. 178 (holotype dorsal), 241 (♂ antennae flagellum), 321 (♂ gen.), 404 (flight period), 427 (distribution)); holotype ♂, Chile, Valdivia, Valdivia, 16. I. 1959, KRAEMER leg.; GP ESN 2633; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

Callipielus perforata NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 81; figs. 172 (holotype dorsal), 173 (♀ dorsal), 237 (♂ antennae flagellum), 313–314 (♂ gen.), 353a (♀ gen.), 377 (bursa copulatrix), 400 (flight period), 426 (distribution)); holotype ♂, Argentina, Rio Negro, S of San Carlos de Bariloche, Pampa del Toro, 900 m, 22.–23. X. 1981, NIELSEN & KARSHOLT leg., sta. 38; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).
Callipielus perforatus [sic]: PASTRANA (2004: 5). — GREHAN (2010: 46).

Callipielus salasi ROBINSON, 1977

ROBINSON (1977: 117; pl. 1, fig. 6 (♂ dorsal), pl. 3, fig. 15 (♂ holotype gen.), pl. 4, fig. 23 (holotype caudal sclerites), pl. 6, figs. 33 (antennal segments), 37 (labium, labial palpus)); holotype ♂, Chile, Cautin, Temuco, Carillanca Experimental Station, 20. V. 1975, SALAS leg.; gen. slide no. 13160; BMNH. — NIELSEN & ROBINSON (1983: 17, 80; figs. 169–170 (♂ dorsal), 171 (♀ dorsal), 235 (♂ antennae flagellum), 236 (♀ antennae flagellum), 266 (♀ foretibia), 311–312 (♂ gen.), 352 (♀ gen.), 376 (bursa copulatrix), 399 (flight period), 425 (distribution)). — ROBINSON & NIELSEN (1984: 16). — GIGANTI et al. (1994: 69). — NIELSEN et al. (2000: 840). — DAPOTO et al. (2003: 100). — PASTRANA (2004: 5).

Callipielus sp.: ROBINSON (1977: 119; pl. 2, fig. 9, ♀ dorsal, pl. 5, fig. 27, ♀ gen., pl. 6, fig. 36, antennal segments); see NIELSEN & ROBINSON (1983: 80).

Callipielus vulgaris NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 87; figs. 182–184 (♂ dorsal), 185–186 (♀ dorsal), 243 (♂ antennae flagellum), 267 (♂ foretibia), 322 (holotype ♂ gen.), 354a–b (♀ gen.), 379 (bursa copulatrix), 406 (flight period), 428 (distribution)); holotype ♂, Argentina, Chubut, Esquel, Lago Menendez, El Sagrario Puerto, 600 m, 2.–4. I. 1982, NIELSEN & KARSHOLT leg., sta. 50; GP ESN 2629; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 66; figs. 1–5, scales). — PASTRANA (2004: 5).

Cibyra WALKER, 1856

Type-species: *Cibyra ferruginosa* WALKER, 1856, by monotypy.

WALKER (1856: 1770). — KIRBY (1892: 938). — QUAIL (1900: 426); *Cibyra sylvinus* [sic] [probably *Triodia sylvina* (LINNAEUS, 1761)]. — NEAVE (1939: 734). — PACLT (1944: 142), syn.: *Alphus* WALLENGREN, 1869, praecoc.: *Alphus* WHITE, 1855 and *Alphus* THOMSON, 1860 [Coleoptera]. — VIETTE (1949c: 102). — PACLT (1953: 145); ◇ (1957: 52). — NYE & FLETCHER (1991: 69). — NIELSEN et al. (2000: 841). — GREHAN & RAWLINS

(2003: 734), syn.: *Aepytyus* HERRICH-SCHÄFFER, [1858]. — GREHAN (2010: 43).

Hepialus (Cibyra): QUAIL (1903: 502), *Hepialus (Cibyra) sylvinus* [sic], probably *Triodia sylvina* (LINNAEUS, 1761).

= *Aepytyus (Xytrops)* VIETTE (1951a: 1); type-species: *Aepytyus (Xytrops) monoargenteus* VIETTE, 1951, by original designation. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17); syn. n.

Xytrops: VIETTE (1951e: 1277). — PACLT (1953: 145); as synonym of *Roseala* VIETTE, 1952. — EDWARDS & HOPWOOD (1966: 315). — NYE & FLETCHER (1991: 323).

Aepytyus (Cibyra): NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Xytrops): NIELSEN et al. (2000: 842).

Remarks. The holotype and a large series of *Aepytyus (Xytrops) monoargenteus* VIETTE, 1951 (type-species) were examined. Morphologically, it is clearly closely related to the type-species of *Cibyra*, which was also examined.

Cibyra danieli (VIETTE, 1961)

Aepytyus danieli VIETTE (1961: 2; fig. 2, [holotype] gen.); holotype ♂, Argentina, Jujuy, Yala, 1450 m, 20. II. 1955, Juan FOERSTER leg.; GP P. VIETTE no. 3789; ZSBS.

Aepytyus (Aepytyus) danieli: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Aepytyus) danieli: NIELSEN et al. (2000: 842).

Cibyra dorita SCHAUS, 1901

SCHAUS (1901: 76); [♂], [Brazil], Paraná, Castro; [GP VIETTE no. 91516, type no. 18607]; USNM.

= *Cibyra poltrona* SCHAUS (1901: 77); [♀], [Brazil], Paraná, Castro; [GP P. VIETTE no. 91515, type no. 18605]; USNM; syn. n.

= *Aepytyus helga* SCHAUS (1929: 55; fig. 22, ♀ dorsal); [holo] type ♀, Brazil, Santa Catarina, [II. 1922, E. D. JONES leg.]; [GP P. VIETTE 91522], type no. 33544; USNM; syn. n.

Hepialus (Cibyra) dorita: PFITZNER (1937: 1292).

Hepialus (Hepialus) helga: PFITZNER (1937: 1291).

Hepialus (Cibyra) poltrona: PFITZNER (1937: 1293).

Xytrops dorita: VIETTE (1951e: 1277).

Aepytyus (Xytrops) dorita: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Aepytyus (Aepytyus) helga: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Aepytyus (Cibyra) poltrona: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Xytrops) dorita: NIELSEN et al. (2000: 842).

Cibyra (Aepytyus) helga: NIELSEN et al. (2000: 842).

Cibyra (Cibyra) poltrona: NIELSEN et al. (2000: 843).

Remarks. The holo- and lectotypes (here designated) of *Aepytyus helga* SCHAUS, 1929, *Cibyra poltrona* SCHAUS, 1901 and *Cibyra dorita* SCHAUS, 1901 were examined and all are conspecific. Matching was done through morphology and also through mtDNA CO-I barcode sequences. One syntype ♂ of *Cibyra dorita* that bears a label “type” is here designated lectotype with the following labels: “Castro, Parana/ *Cibyra dorita* type SCHAUS/ Type n°. 18607 USNM/ Genitalia slide P. VIETTE USNM 91516/ Photo, det. E. S. NIELSEN 1984”. One syntype ♀ of *Cibyra poltrona* that bears label “type” is here designated lectotype with the following labels: “Castro, Parana/ *Cibyra poltrona* type SCHAUS/ Type n°. 18605 USNM/ Genitalia slide P. VIETTE USNM 91515/ Photo, det. E. S. NIELSEN 1984”. The two designations are made to permanently stabilise the species’ identification.

***Cibyra ferruginosa* WALKER, 1856**

WALKER (1856: 1770), Brazil, coll. STEVEN; BMNH. — KIRBY (1892: 938). — VIETTE (1949c: 102); ♂ (1951c: 95), syn.: *Cibyra dormita* SCHAUS, 1901; ♂ (1951e: 1277). — GREHAN (2010: 45).

Cibyra ferruginea [sic]: KIRBY (1892: 866).

= *Cibyra dormita* SCHAUS (1901: 77), [Brazil, Rio de Janeiro], Petrópolis; [coll. SCHAUS]; [type no. 18602]; USNM. — VIETTE (1950a: 75; fig. 1 ♂ gen.), cited as *Aepytus exclamans* (HERRICH-SCHÄFFER, [1854]), see VIETTE (1951c: 95); ♂ (1951c: 95; fig. 2 [syn-]type ♂ gen.), as synonym of *Cibyra ferruginosa* WALKER, 1856.

Hepialus (Cibyra) dormita: PFITZNER (1937: 1293).

Aepytus (Cibyra) dormita: NIELSEN & ROBINSON (1983: 20); as synonym of *Aepytus (Cibyra) ferruginosa* (WALKER, 1856). — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytus (Cibyra) ferruginosa* (WALKER, 1856).

Aepytus (Cibyra) ferruginosa: NIELSEN & ROBINSON (1983: 20), syn.: *Aepytus (Cibyra) dormita* (SCHAUS, 1901). — ROBINSON & NIELSEN (1984: 17), syn.: *Aepytus (Cibyra) dormita* (SCHAUS, 1901).

Cibyra (Cibyra) dormita: NIELSEN et al. (2000: 843), as synonym of *Cibyra (Cibyra) ferruginosa* WALKER, 1856.

Cibyra (Cibyra) ferruginosa: NIELSEN et al. (2000: 843), syn.: *Cibyra (Cibyra) dormita* SCHAUS, 1901. — GREHAN (2010: 51; fig. c, appendix).

***Cibyra forsteri* (VIETTE, 1961)**

Aepytus forsteri VIETTE (1961: 1; fig. 1, [holotype] gen.); holotype ♂, Bolivia, Cochabamba, 2600 m, 10. XI. 1956, R. ZISCHKA leg.; GP P. VIETTE no. 3552; ZSBS.

Aepytus (Aepytus) forsteri: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Aepytus) forsteri: NIELSEN et al. (2000: 842).

***Cibyra monoargenteus* (VIETTE, 1951)**

Aepytus (Xytrops) monoargenteus VIETTE (1951a: 2; fig. 3 [holotype] ♂ gen.); [holo-]type ♂, Brazil, Paraná, Curitiba [RECTE Curitiba], February; GP P. VIETTE no. 1345; MNHN. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Xytrops) monoargenteus: NIELSEN et al. (2000: 842). — GREHAN (2010: 54; fig. n, appendix).

***Cibyra munona* (SCHAUS, 1929)**

Aepytus munona SCHAUS (1929: 56; fig. 21, ♂ dorsal); [holo-]type ♂, Brazil, Santa Catarina, [ii. 1922, E. D. JONES leg.]; [GP P. VIETTE no. 91521], type no. 33545; USNM.

Hepialus (Hepialus) munona: PFITZNER (1937: 1291).

Aepytus (Aepytus) munona: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Aepytus) munona: NIELSEN et al. (2000: 842).

***Cibyra oreas* (SCHAUS, 1892), comb. rev.**

Dalaca oreas SCHAUS (1892: 330); Brazil, [Rio de Janeiro], Petrópolis, SCHAUS leg.; coll. SCHAUS; [GP P. VIETTE no. 91527, type no. 18604]; USNM. — BERTRAU (1893: 190). — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1295).

Aepytus oreas: SCHAUS (1929: 56).

Aepytus (Paragorgopis) oreas: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopis) oreas: NIELSEN et al. (2000: 842).

Vietteogorgopis oreas: ÖZDIKMEN (2007: 117).

Remarks. The lectotype (here designated) was examined, and through morphology it is clearly related to the type-species of the genus *Cibyra* WALKER, 1856. — One syntype ♀ of *Cibyra oreas*

(SCHAUS, 1892) that bears a label “type” is here designated lectotype with the following labels: /Petrópolis, Brazil/ *Cibyra oreas* type SCHAUS/ Coll. Wm. SCHAUS/ Type n°. 18604 USNM/ Genitalia slide P. VIETTE USNM 91527/ Photograph on file USNM/ Photo, det. E. S. NIELSEN 1984/. The designation is made to permanently stabilise the species' identification.

***Cibyra pluriargenteus* (VIETTE, 1956)**

Xytrops pluriargenteus VIETTE (1956: 378; fig. 4, ♂ gen.); holotype ♂, Brazil, São Paulo, Alto da Serra, i. 1923, R. SPITZ leg.; GP P. VIETTE no. 2296; BMNH.

Aepytus (Xytrops) pluriargenteus: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Xytrops) pluriargenteus: NIELSEN et al. (2000: 843).

***Cibyra schausi* (VIETTE, 1952), comb. rev.**

Paragorgopis schausi VIETTE (1952a: 142; fig. 3, ♀ gen.); holotype ♀, Brazil, São Paulo, Araras, J. G. FOETTERLE [leg.]; GP P. VIETTE no. 2260; NHMW.

Aepytus (Paragorgopis) schausi: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopis) schausi: NIELSEN et al. (2000: 842).

Vietteogorgopis schausi: ÖZDIKMEN (2007: 117). — KOÇAK & KOÇAK (2008: 31).

Remarks. The holotype was examined, and through morphology it is clearly related to the type-species of the genus *Cibyra* WALKER, 1856.

***Cibyra stigmatica* (PFITZNER, 1937), comb. n.**

Dalaca stigmatica PFITZNER (1937: 1296; pl. 100b, [♀] dorsal); no data; coll. SEITZ; [SMFL]. — ZUKOWSKI (1954: 93); as synonym of *Dalaca tesselloides* SCHAUS, 1901. — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. Based on large series of this taxon and ♂ genitalia, it is clearly closely related to the type-species of the genus *Cibyra* WALKER, 1856. One syntype ♀ deposited in SMFL of *Dalaca stigmatica* PFITZNER, 1937 is here designated lectotype with the following labels: /stigmatica, S^a Catarina (?) od. Costarica/ Coll. A. SEITZ. The designation is made to permanently stabilise the species' identification.

***Cibyra verresi* (SCHAUS, 1929)**

Aepytus verresi SCHAUS (1929: 56; fig. 23, ♂ dorsal); [holo-]type ♂, Brazil, Santa Catarina, [2. IV. 1924, E. D. JONES leg.]; [GP P. VIETTE no. 91523], type no. 33546; USNM.

Dalaca verresi: PFITZNER (1937: 1296).

Aepytus (Xytrops) verresi: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Xytrops) verresi: NIELSEN et al. (2000: 843).

***Cibyra yungas* (VIETTE, 1961)**

Xytrops yungas VIETTE (1961: 3; fig. 4, [holotype] ♂ gen.); holotype ♂, Bolivia, [Cochabamba], Yungas del Palmar, 2000 m, R. ZISCHKA leg.; GP P. VIETTE no. 3790; ZSBS.

Aepytus (Xytrops) yungas: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Xytrops) yungas: NIELSEN et al. (2000: 842).

***Cibyra zischkai* (VIETTE, 1961)**

Aepytus zischkai VIETTE (1961: 2; fig. 3, [holotype] ♂ gen.); holotype ♂, Bolivia, Cochabamba, 2600 m, 5. XII. 1954, R. ZISCHKA leg.; GP P. VIETTE no. 3788; ZSBS.

Aepytus (Aepytus) zischkai: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Aepytus) zischkai: NIELSEN et al. (2000: 842).

Dalaca WALKER, 1856

Type-species: *Dalaca nigricornis* WALKER, 1856 by subsequent designation by DRUCE (1887: 232).

Dalaca WALKER (1856: 1549, 1559); included species: *nigricornis* WALKER, 1856, *exul* (HERRICH-SCHÄFFER, [1853]), *nomaqua* WALKER, 1856, *exclamans?* (HERRICH-SCHÄFFER, [1854]), *epigramma?* (HERRICH-SCHÄFFER, [1854]).

GERSTAECKER (1857: 425). — HERRICH-SCHÄFFER ([1858]: 56; as synonym of *Triodia* HÜBNER, [1820], *Aepytus* HERRICH-SCHÄFFER, [1856]. — BUTLER (1882: 25). — DRUCE (1887: 232). — KIRBY (1892: 886); type-species: *Dalaca nomaqua* WALKER 1856, by subsequent designation [incorrect designation]. — DRUCE (1898: 450). — QUAIL (1900: 432). — WAGNER & PFITZNER (1911: 13). — STRAND (1914: 59). — JANSE (1919: 240). — GAEDE (1930: 555); misidentification. — PFITZNER (1937: 1293). — NEAVE (1939: 8). — JANSE (1942: 5). — PAULT (1944: 143); ◇ (1949: 152), syn.: *Huapina* BRYK, 1944. — VIETTE (1950: 144), syn.: *Huapina* BRYK, 1944, *Maculella* VIETTE, 1950; ◇ (1951d: 74, 80). — PAULT (1953: 143), syn.: *Blanchardina* VIETTE, 1950. — ZUKOWSKI (1954: 94). — DUMBLETON (1966: 924). — SCHRÖDER (1967: 339). — NIELSEN & ROBINSON (1983: 16, 51), syn.: *Huapina* BRYK, 1944, *Maculella* VIETTE, 1950, *Toenga* TINDALE, 1954. — MALLET (1984: 77). — ROBINSON & NIELSEN (1984: 16), syn.: *Huapina* BRYK, 1944, *Maculella* VIETTE, 1950, *Toenga* TINDALE, 1954. — NIELSEN & SCOBLE (1986: 43). — GREHAN (1989: 805). — NYE & FLETCHER (1991: 87). — KRISTENSEN (1998: 62). — NIELSEN et al. (2000: 839), syn.: *Huapina* BRYK, 1944, *Maculella* VIETTE, 1950, *Toenga* TINDALE, 1954. — SIMONSEN (2002: 65). — GREHAN (2010: 49; fig. L, appendix).

= *Huapina* BRYK (1944: 28); type-species: *Huapina parvigguttata* BRYK, 1944, by monotypy. — PAULT (1949: 152); as synonym of *Dalaca* WALKER, 1856. — NEAVE (1950: 119). — VIETTE (1950e: 144); as synonym of *Dalaca* WALKER, 1856; ◇ (1951d: 80). — NIELSEN & ROBINSON (1983: 16); as synonym of *Dalaca* WALKER, 1856. — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca* WALKER, 1856. — NYE & FLETCHER (1991: 153). — NIELSEN et al. (2000: 839); as synonym of *Dalaca* WALKER, 1856.

= *Maculella* VIETTE (1950c: 55); type-species: *Dalaca noctuides* PFITZNER, 1914 by original designation. — VIETTE (1950e: 144); as synonym of *Dalaca* WALKER, 1856; ◇ (1951d: 76). — EDWARDS & HOPWOOD (1966: 157). — SCHRÖDER (1967: 338). — NIELSEN & ROBINSON (1983: 16); as synonym of *Dalaca* WALKER, 1856. — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca* WALKER, 1856. — NYE & FLETCHER (1991: 182). — NIELSEN et al. (2000: 839); as synonym of *Dalaca* WALKER, 1856.

= *Toenga* TINDALE (1954: 13); type-species: *Toenga oceanica* TINDALE, 1954, by original designation. — PAULT (1957: 52). — DUMBLETON (1966: 924, 971). — EDWARDS & HOPWOOD (1966: 296). — NIELSEN & ROBINSON (1983: 16, 51); as synonym of *Dalaca* WALKER, 1856. — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca* WALKER, 1856. — NYE & FLETCHER (1991: 309). — NIELSEN et al. (2000: 839); as synonym of *Dalaca* WALKER, 1856.

Dalaca chiliensis (VIETTE, 1950)

Maculella chiliensis VIETTE (1950c: 57; fig. 6, ♂ gen.); [holo-]type ♂, Chile, Valdivia, Arturo von LOSSBERG [leg.], 1901; coll. BIEDERMANN, ex coll. OBERTHÜR; GP P. VIETTE no. 950; MNHN. — DURÁN (1976: 121, 123).

Dalaca chilensis [sic]: VIETTE (1950c: 55, 56).

Meculella [sic] *chiliensis*: CARRILLO (1974: 46).

Dalaca chiliensis: NIELSEN & ROBINSON (1983: 16, 56; figs. 45 (ductus bursae), 77–80 (♂ dorsal), 81 (♀ dorsal), 215 (♂ antennae flagellum), 216 (♀ antennae flagellum), 254 (♂ fore-

tibia), 255 (♀ foretibia), 280–281 (♂ gen.), 335 (♀ gen.), 365 (bursa copulatrix), 388 (flight period), 418 (distribution)). — ROBINSON & NIELSEN (1984: 16). — CISTERNAS (1989: 10); ◇ (1992: 88); ◇ (2000: 1). — KOCH & WATERHOUSE (2000: 31, 72, 87, 121, 142, 159). — NIELSEN et al. (2000: 839). — JACKSON (2007: 35). — CISTERNAS et al. (2007: 42). — PAPE (2009: 4).

Dalaca crocatus (URETA, 1956)

Hepialus crocatus URETA (1956: 284); holotype ♂, Chile, “Araucanía [Cautin and Malleco], ii. 1892”; coll. M.N. no. 2247; [GP ESN 2634, type no. 759]; MNHC. — URETA (1957; pl. 1, fig. 8, ♂ dorsal). — CAMOUSSEIGHT (1980: 31).

Dalaca crocatus: NIELSEN & ROBINSON (1983: 16, 55; figs. 76 (♂ holotype), 214 (antennae flagellum), 279 (♂ gen.), 387 (flight period), 418 (distribution)). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839).

Dalaca laminata NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 67; figs. 129–131 (♂ dorsal), 132 (♀ dorsal), 224 (♂ antennae flagellum), 225 (♀ antennae flagellum), 260 (♀ foretibia), 293 (♂ gen.), 294 (holotype gen.), 343–344 (♀ gen.), 369 (bursa copulatrix), 393 (flight period), 421 (distribution)); holotype ♂, Chile, Malleco, Cordillera Nahuelbuta, Cabrería, 9.-15. i. 1977, PEÑA [leg.]; GP ESN 2561; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839).

Dalaca nigricornis WALKER, 1856

WALKER (1856: 1560); ♂, Chile, coll. CUMING; BMNH. — BUTLER (1882: 25); as synonym of *Dalaca venosa* (BLANCHARD, 1852). — AURIVILLIUS (1884: 524); as synonym of *Dalaca venosa* (BLANCHARD, 1852). — DRUCE (1887: 232). — KIRBY (1892: 886); as synonym of *Dalaca venosa* (BLANCHARD, 1852). — WAGNER & PFITZNER (1911: 14); as synonym of *Dalaca venosa* (BLANCHARD, 1852). — PFITZNER (1937: 1295); as synonym of *Dalaca venosa* (BLANCHARD, 1852). — VIETTE (1950e: 145; fig. 3 type ♂ gen.). — NIELSEN & ROBINSON (1983: 17, 64; figs. 50 (hindwing apex), 124 (holotype [RECTE syntype] ♂ dorsal), 125–126 (♀ dorsal), 221 (♂ antennae flagellum), 222 (♂ [error, ♀] antennae flagellum), 259 (♀ foretibia), 290 (holotype [RECTE syntype] gen.), 342 (♀ gen.), 368 (bursa copulatrix), 391 (flight period), 420 (distribution)); holotype [RECTE syntype] ♂, Chile; gen. slide no. 2086. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839). — PASTRANA (2004: 5).

Dalaca pallens (BLANCHARD, 1852)

Hepialus pallens BLANCHARD (1852: 69; pl. 4, fig. 5 dorsal); Chile, Coquimbo. — WALKER (1856: 1556). — KIRBY (1892: 884). — WAGNER & PFITZNER (1911: 9). — LLOYD & BLACKMAN (1966: 14). — CARRILLO (1974: 46). — DURÁN (1976: 123).

Dalaca (*Triodia*) *venosa* [sic] [misidentification]; BERG (1882: 219); see NIELSEN & ROBINSON (1983: 90).

= *Aepytus dimidiatus* BERG (1882: 220); single ♂ [holotype], Chile, Concepcion, 26. ii. 1879, BERG leg. — AURIVILLIUS (1884: 524). — KIRBY (1892: 887).

= *Dalaca hemileuca* BUTLER (1882: 27); 1 ♂, 1 ♀, Chile, EDMONDS leg.; BMNH. — AURIVILLIUS (1884: 524). — KIRBY (1892: 887). — WAGNER & PFITZNER (1911: 13). — PFITZNER (1937: 1295). — NIELSEN & ROBINSON (1983: 16, 57); lectotype ♂; as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).

= *Dalaca marmorata* BUTLER (1882: 25); 2 ♂♂, Chile, Las Zonas, EDMONDS leg.; BMNH. — AURIVILLIUS (1884: 524). — KIRBY (1892: 887). — WAGNER & PFITZNER (1911: 14). —

- PFITZNER (1937: 1295). — NIELSEN & ROBINSON (1983: 16, 57); lectotype ♂; GP BMNH Hepial. 10777; as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- = *Dalaca pallens*: BUTLER (1882: 26). — PFITZNER (1937: 1295). — NIELSEN & ROBINSON (1983: 16, 57; figs. 46–47 (♂ and ♀ forewing base), 48 (saccus), 82–96 (♂ dorsal), 97–99 (♀ dorsal), 217 (♂ antennae flagellum), 218 (♀ antennae flagellum), 256 (♂ foretibia), 257 (♀ foretibia), 282–286 (♂ gen.), 336–338 (♀ gen.), 366 (bursa copulatrix), 389 (flight period), 419 (distribution)), syn.: *Dalaca hemileuca* BUTLER, 1882, *Dalaca marmorata* BUTLER, 1882, *Dalaca subfervens* BUTLER, 1882, *Dalaca violacea* BUTLER, 1882, *Dalaca dimidiatus* (BERG, 1882), *Dalaca noctuides* PFITZNER, 1914, *Dalaca parviguttata* (BRYK, 1944), *Dalaca pseudodimiata* (PACLT, 1953), *Dalaca oceanica* (TINDALE, 1954), *Dalaca venosa* [sic] BERG (1882, nec BLANCHARD 1852). — ROBINSON & NIELSEN (1984: 16), syn.: *Dalaca hemileuca* BUTLER, 1882, *Dalaca marmorata* BUTLER, 1882, *Dalaca subfervens* BUTLER, 1882, *Dalaca violacea* BUTLER, 1882, *Dalaca dimidiatus* (BERG, 1882), *Dalaca noctuides* PFITZNER, 1914, *Dalaca parviguttata* (BRYK, 1944), *Dalaca pseudodimiata* (PACLT, 1953), *Dalaca oceanica* (TINDALE, 1954). — CISTERNAS (1989: 10); ♂ (1992: 88). — GIGANTI et al. (1994: 70). — CISTERNAS (2000: 1). — KOCH & WATERHOUSE (2000: 31, 72, 87, 121, 142, 159). — NIELSEN et al. (2000: 839), syn.: *Dalaca hemileuca* BUTLER, 1882, *Dalaca marmorata* BUTLER, 1882, *Dalaca subfervens* BUTLER, 1882, *Dalaca violacea* BUTLER, 1882, *Dalaca dimidiatus* (BERG, 1882), *Dalaca noctuides* PFITZNER, 1914, *Dalaca parviguttata* (BRYK, 1944), *Dalaca pseudodimiata* (PACLT, 1953), *Dalaca oceanica* (TINDALE, 1954). — SIMONSEN (2002: 66; figs. 11–19, scales). — CISTERNAS et al. (2003: 51). — DAPOTO et al. (2003: 100). — PASTRANA (2004: 5), syn.: *Dalaca noctuides* PFITZNER, 1914. — DEVOTTO et al. (2007: 508). — CISTERNAS et al. (2007: 42). — DEVOTTO et al. (2008: 228). — AGUILERA et al. (2009: 35). — PAPE (2009: 4).
- = *Dalaca subfervens* BUTLER (1882: 25); Chile, Las Zonas, EDMONDS leg.; BMNH. — AURIVILLIUS (1884: 524). — KIRBY (1892: 886). — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1295). — NIELSEN & ROBINSON (1983: 16, 57); lectotype ♂; as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- = *Dalaca violacea* BUTLER (1882: 26); 1 ♂ [holotype], Chile, EDMONDS leg.; BMNH. — AURIVILLIUS (1884: 524). — KIRBY (1892: 887). — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1295). — NIELSEN & ROBINSON (1983: 16, 57); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Hepialus dimidiatus*: WAGNER & PFITZNER (1911: 4).
- = *Dalaca noctuides* PFITZNER (1914: 105); Chile, Valdivia; coll. PFITZNER; [SMFL]; ♂ (1937: 1295; pl. 99e [♂] dorsal). — ZUKOWSKI (1954: 94); as synonym of *Dalaca fusca* [sic] MABILLE, 1885. — IHL (1947: 78). — CABALLERO (1955: 7). — ISLA (1959: 16). — CARRILLO (1974: 46). — NIELSEN & ROBINSON (1983: 17, 57); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — PASTRANA (2004: 5); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Hepialus (Hepialus) dimidiatus*: PFITZNER (1937: 1291).
- = *Huapina parviguttata* BRYK (1944: 28; pl. 2, fig. 17 (♀ dorsal)); [holo-]type ♀, [Argentina, Rio Negro, San Carlos de Bariloche], Nahuel Huapí, Península Llau Llau, N Puerto Nuevo, Patagonia, 770–780 m, x. 1933–III. 1934, LJUNGNER leg.; NRSS.
- Dalaca parviguttata*: PACLT (1949: 149, 152); as synonym of *Dalaca venosa* [sic] (BLANCHARD). — VIETTE (1950e: 145; figs. 1–2 [holo-]type ♀ gen.); GP P. VIETTE no. 2171. — PACLT (1953: 146); as synonym of *Dalaca venosa* [sic] (BLANCHARD). — NIELSEN & ROBINSON (1983: 17, 57); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Aepytus dimidiatus* [misidentification]: PACLT (1949: 149); see PACLT (1953: 145).
- Maculella hemileuca*: VIETTE (1950c: 55); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Maculella pallens*: VIETTE (1950c: 55).
- Maculella marmorata*: VIETTE (1950c: 55); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Maculella noctuides*: VIETTE (1950c: 56; fig. 5, ♂ gen.). — SCHRÖDER (1967: 338); lectotype ♂, SMFT 1002; GP P. VIETTE no. 1410. — DURÁN (1976: 121, 123). — RODRÍGUEZ et al. (1980: 73).
- Maculella subfervens*: VIETTE (1950c: 55); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Maculella violacea*: VIETTE (1950c: 55); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- = *Lossbergiana pseudodimiata* PACLT (1953: 145); [holo-]type ♀, [Argentina, Rio Negro, San Carlos de Bariloche], Nahuel Huapí, Península Llau Llau, N of Puerto Nuevo, Patagonia, 770–780 m, x. 1933–III. 1934, LJUNGNER leg., syn.: *Aepytus dimidiatus* PACLT, 1949, nec BERG.
- = *Toenga oceanica* TINDALE (1954: 15; figs. 1a (antenna), 1b (labial palpus), 1c (venation), 1d–1e (abdomen), pl. 1, fig. [2] ([holotype] ♀ dorsal)); single specimen [= holotype ♀], Cook Islands, Rarotonga [mislabelled, probably Chile or Argentina]; no. 93: 162; BMNH. — TINDALE (1981: 966).
- Dalaca dimidiatus*: VIETTE (1961: 7). — CARRILLO (1974: 46). — NIELSEN & ROBINSON (1983: 17, 57); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Maculella (Dalaca) noctuides*: LLOYD & BLACKMAN (1966: 14).
- Maculella dimidiata* [sic]: DURÁN (1976: 119, 127).
- Dalaca pseudodimiata*: NIELSEN & ROBINSON (1983: 17, 57); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Dalaca oceanica*: NIELSEN & ROBINSON (1983: 17, 57); holotype GP BMNH Hepial. 20444; as synonym of *Dalaca pallens* (BLANCHARD, 1852). — ROBINSON & NIELSEN (1984: 16); as synonym of *Dalaca pallens* (BLANCHARD, 1852). — NIELSEN et al. (2000: 839); as synonym of *Dalaca pallens* (BLANCHARD, 1852).
- Dalaca dimidiata* [sic]: PASTRANA (2004: 5).
- Dalaca parafuscus* NIELSEN, ROBINSON & WAGNER, 2000**
- Hepialus fuscus* MABILLE (1885: 56); [Chile, Magallanes], “Ex insulis Magellanicas”; *praeocc.*: *Hepialus fuscus* HAWORTH, 1809 [Hepialidae]. — MABILLE (1888: 8; fig. 6 dorsal). — STAUDINGER (1899: 44). — PAGENSTECHER (1902: 398, 399). —

ENDERLEIN (1912: 90). — VIETTE & FLETCHER (1968: 392); type not found.

Dalaca fusca [sic]: KIRBY (1892: 887). — WAGNER & PFITZNER (1911: 13). — PFITZNER (1937: 1295; pl. 185b, [♂] dorsal). — ZUKOWSKI (1954: 94), syn.: *Dalaca noctuides* PFITZNER, 1914. — PASTRANA (2004: 5).

Dalaca fuscus: NIELSEN & ROBINSON (1983: 17, 68; figs. 133–135 (♂ dorsal), 136 (♀ dorsal), 226 (♂ antennae flagellum), 227 (♀ antennae flagellum), 295–296 (♂ gen.), 345 (♀ gen.), 370 (bursa copulatrix), 394 (flight period), 422 (distribution)). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839). = *Dalaca parafuscus* NIELSEN, ROBINSON & WAGNER (2000: 839); replacement name for *Hepialus fuscus* MABILLE, 1885.

Dalaca patriciae NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 66; figs. 127 (holotype ♂ dorsal), 128 (♂ dorsal), 223 (♂ antennae flagellum), 291–292 (♂ gen.), 392 (flight period), 421 (distribution)); holotype ♂, Argentina, Neuquen, San Martin de los Andes, Cerro Chapelco, 1400–1650 m, 1. XII. 1981, NIELSEN & KARSHOLT leg., sta. 36; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839). — PASTRANA (2004: 5).

Dalaca postvariabilis NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 69; figs. 137–141 (♂ dorsal), 142–143 (♀ dorsal), 228 (♂ antennae flagellum), 229 (♀ antennae flagellum), 261 (♀ foretibia), 297–298 (♂ gen.), 346 (♀ gen.), 371 (bursa copulatrix), 395 (flight period), 422 (distribution)); holotype ♂, Argentina, Neuquen, Lago Lacar, Pucara, 650 m, 26. XII. 1978, Mision Cientifica Danesa leg., sta. 9; GP ESN 2543; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839). — SIMONSEN (2002: 66; figs. 20–23). — PASTRANA (2004: 5).

Dalaca quadricornis NIELSEN & ROBINSON, 1983

NIELSEN & ROBINSON (1983: 17, 62; figs. 49 (saccus), 100–119 (♂ dorsal), 120–123 (♀ dorsal), 219 (♂ antennae flagellum), 220 (♀ antennae flagellum), 258 (♀ foretibia), 287–289 (♂ gen.), 339–341 (♀ gen.), 367 (bursa copulatrix), 390 (flight period), 420 (distribution)); holotype ♂, Argentina, Chubut, Esquel, Lago Menendez, El Sagrario Puerto, sta. 50, 600 m, 2.–4. I. 1982, NIELSEN & KARSHOLT leg.; GP ESN 2834; ZMUC. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 839). — PASTRANA (2004: 5).

Dalaca variabilis (VIETTE, 1950)

Maculella variabilis VIETTE (1950c: 57; fig. 7, ♂ gen.); [holo-]type ♂, Chile, Valdivia, Arturo von LOSSBERG [leg.], 1901; coll. BIEDERMANN, ex coll. OBERTHÜR; GP P. VIETTE no. 946; MNHN.

Dalaca variabilis: NIELSEN & ROBINSON (1983: 17, 71; figs. 144–154 (♂ dorsal), 155–158 (♀ dorsal), 230 (♂ antennae flagellum), 231 (♀ antennae flagellum), 299–302 (♂ gen.), 347–348 (♀ gen.), 372–373 (bursa copulatrix), 396 (flight period), 423 (distribution)). — ROBINSON & NIELSEN (1984: 16). — CISTERNAS (2000: 1). — NIELSEN et al. (2000: 839). — CISTERNAS et al. (2007: 42).

Dalaca variables [sic]: CISTERNAS (2000: 4).

Druceiella VIETTE, 1949

Type-species: *Hepialus momus* DRUCE, 1890, by original designation.

Druceiella VIETTE (1949a: 52); included species: *metellus* (DRUCE, 1890), *momus* (DRUCE, 1890).

VIETTE (1951d: 75). — PAULT (1953: 145); as synonym of *Pseudophassus* PFITZNER & GAEDE [RECTE *Pseudophassus*

WEYMER [ms.] in PFITZNER 1938). — EDWARDS & HOPWOOD (1966: 83). — SCHRÖDER (1967: 342). — NIELSEN & ROBINSON (1983: 18, 111). — ROBINSON & NIELSEN (1984: 16). — NYE & FLETCHER (1991: 100). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN (2010: 43).

Druceiella amazonensis VIETTE, 1950

VIETTE (1950d: 168; figs. 10 (holotype 8th tergite), 11 (holotype gen.)); holotype ♂, Brazil, [Pará], Óbidos; GP P. VIETTE no. 2113; MNHN. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

Druceiella basirubra (SCHAUS, 1901)

Dalaca basirubra SCHAUS (1901: 76); Peru; [GP VIETTE no. 2628]; USNM. — WAGNER & PFITZNER (1911: 13). — PFITZNER (1937: 1296). — ZUKOWSKI (1954: 93).

= *Pseudophassus metricus* var. *songoensis* PFITZNER (1914: 110); Bolivia, Rio Songo, [1912, FASSL leg.]; coll. PFITZNER; [SMFL]. — FORBES (1942: 406); as synonym of *Dalaca metellus* (DRUCE, 1890). — VIETTE (1961: 6); as synonym of *Druceiella basirubra* (SCHAUS, 1901).

Pseudophassus momus songoensis: PFITZNER (1938: 1301).

Dalaca metricus var. *songoensis*: FORBES (1942: 406); as synonym of *Dalaca metellus* (DRUCE, 1890).

Druceiella metellus [misidentification]: VIETTE (1949a: 54; figs. 4 (8th ♂ tergite), 5 (♂ gen.), 9 (8th sternite)); see VIETTE (1950d: 167), NIELSEN & ROBINSON (1983: 112).

Druceiella songoensis: VIETTE (1950d: 167; figs. 4–9, 8th ♂ tergite). — SCHRÖDER (1967: 342); lectotype [♂], SMFT 1013. — NIELSEN & ROBINSON (1983: 18); as synonym of *Druceiella basirubra* (SCHAUS, 1901). — ROBINSON & NIELSEN (1984: 16); as synonym of *Druceiella basirubra* (SCHAUS, 1901). — NIELSEN et al. (2000: 840); as synonym of *Druceiella basirubra* (SCHAUS, 1901).

Druceiella basirubra: VIETTE (1961: 6), syn.: *Pseudophassus metricus* var. *songoensis* PFITZNER, 1914. — NIELSEN & ROBINSON (1983: 18, 112; figs. 74a (prelabium, labial palpus), 74b (venation), 75 (8th sternite, tergite), 213 (♂ dorsal), 253 (♂ antennae flagellum), 278 (♂ foretibia), 334 (♂ gen.), 364 (♀ gen.), 386 (bursa copulatrix), 416 (flight period), 431 (distribution)), syn.: *Druceiella songoensis* (PFITZNER, 1914). — ROBINSON & NIELSEN (1984: 16), syn.: *Druceiella songoensis* (PFITZNER, 1914). — NIELSEN et al. (2000: 840), syn.: *Druceiella songoensis* (PFITZNER, 1914). — SIMONSEN (2002: 65). — GREHAN (2010: 45).

Pseudophassus monus [sic] *songoensis*: SCHRÖDER (1967: 342).

Druceiella metellus (DRUCE, 1890)

Hepialus metellus DRUCE (1890: 509; fig. 2 [♂] dorsal); Ecuador, Sarayacu, BUCKLEY leg.; coll. DRUCE; [GP P. VIETTE, BMNH GP Hepial. 2081]; [BMNH]. — BERTKAU (1891: 196).

Phassus metellus: KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 18).

Dalaca metellus: HAMPSON (1903: 260). — FORBES (1942: 406), syn.: *Hepialus momus* DYAR, 1915, nec DRUCE, 1890, *Pseudophassus metricus*, var. *songoensis* PFITZNER 1914, var. [sic] *momus metricus* PFITZNER, 1938.

Hepialus momus [misidentification]: DYAR (1915b: 350); see FORBES (1942: 406).

Pseudophassus momus metellus: PFITZNER (1938: 1301; pl. 99g, [♂] dorsal). — FORBES (1942: 406); as synonym of *Dalaca metellus*.

Druceiella metellus: VIETTE (1950d: 167; figs. 1 8th ♂ tergite, 2 ♂ gen.); ◇ (1961: 6). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

***Druceiella momus* (DRUCE, 1890)**

Hepialus momus DRUCE (1890: 508; fig. 3 [♂] dorsal); Ecuador, Sarayacu, BUCKLEY leg.; coll. DRUCE; [GP P. VIETTE no. BM Gen slide 2085]; [BMNH]. — BERKKAU (1891: 196).

Phassus momus: KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 18).

Pseudophassus momus: PFITZNER (1938: 1301; pl. 185e [♂] dorsal). — ZUKOWSKI (1954: 94).

= *Pseudophassus momus* f. *metricus* PFITZNER (1938: 1301; pl. 99h [♂] dorsal); Bolivia, Rio Songo, FASSL [leg.].

Pseudophassus var. [sic] *metricus*: FORBES (1942: 406), as synonym of *Dalaca metellus* (DRUCE, 1890).

Druceiella momus: VIETTE (1949a: 53; figs. 6 (8th ♂ tergite), 7 (8th ♂ sternite), 8 (♂ gen.)). — VIETTE (1961: 6). — ROBINSON & NIELSEN (1984: 16), syn.: *Druceiella metricus* (PFITZNER, 1938). — NIELSEN et al. (2000: 840), syn.: *Druceiella metricus* (PFITZNER, 1938).

Druceiella nomus [sic]: NIELSEN & ROBINSON (1983: 18), syn.: *Druceiella metricus* (PFITZNER, 1938).

Druceiella metricus: NIELSEN & ROBINSON (1983: 18); as synonym of *Druceiella momus* (DRUCE, 1890). — ROBINSON & NIELSEN (1984: 16); as synonym of *Druceiella momus* (DRUCE, 1890). — NIELSEN et al. (2000: 840); as synonym of *Druceiella momus* (DRUCE, 1890).

***Gymelloxes* VIETTE, 1952, stat. rev.**

Type-species: *Dalaca terea* SCHAUS, 1892, by original designation.

Gymelloxes VIETTE (1952b: 27); included species: *terea* (SCHAUS, 1892), *trilinearis* (PFITZNER, 1914).

PAULT (1957: 51); as synonym of *Aepytyus* HERRICH-SCHÄFFER, [1858]. — EDWARDS & HOPWOOD (1966: 114). — NYE & FLETCHER (1991: 139).

Aepytyus (*Gymelloxes*): NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Gymelloxes*): NIELSEN et al. (2000: 842). — GREHAN (2010: 44).

Remarks. *Gymelloxes* VIETTE, 1952 was described mainly based on the ♂ genitalia characters, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

***Gymelloxes prosopus* (DRUCE, 1901), comb. n.**

Hepialus prosopus DRUCE (1901: 436); Colombia, Bonda; coll. DRUCE; BMNH.

Hepialus propopus [sic]: WAGNER & PFITZNER (1911: 9).

= *Dalaca chiriquensis* PFITZNER (1914: 105); [Panama], Chiriqui; coll. PFITZNER; [SMFL]; ♂ (1937: 1294; pl. 99b [♀] dorsal). — SCHRÖDER (1967: 339); lectotype [♀], SMFT 81. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 843); **syn. n.**

= *Dalaca muysca* PFITZNER (1914: 105); [Panama], Chiriqui; coll. PFITZNER; [SMFL]. — SCHRÖDER (1967: 340); “holotype” [RECTE lectotype] [♀], SMFT 99, syn.: *Dalaca terea* SCHAUS, 1892; **syn. n.**

Dalaca terea f. *muysca*: PFITZNER (1937: 1294; pl. 99f [♀] dorsal).

Pseudophassus prosopus PFITZNER (1938: 1301).

Aepytyus (*Gymelloxes*) *muysca*: NIELSEN & ROBINSON (1983: 19); as synonym of *Aepytyus* (*Gymelloxes*) *terea* (SCHAUS, 1892). — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytyus* (*Gymelloxes*) *terea* (SCHAUS, 1892).

Pfütznieriana prosopus: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

Cibyra (*Gymelloxes*) *muysca*: NIELSEN et al. (2000: 842); as synonym of *Cibyra* (*Gymelloxes*) *terea* (SCHAUS, 1892).

Remarks. The lectotypes of *Dalaca chiriquensis* PFITZNER, 1914, *Dalaca muysca* PFITZNER, 1914 and *Hepialus prosopus* DRUCE, 1901 (here designated) were examined and all are conspecific. In the BMNH, there is one ♀ of *Gymelloxes prosopus* (DRUCE, 1901) that bears a label “type” which is here designated lectotype; it has the following labels: /*Hepialus prosopus* DRUCE type/ Bonda, Colombia, 150 ft., H. H. SMITH/ Sept/ JOICEY Coll. B.M. 1929-122/ Ex Coll. Herbert DRUCE 1913/. The designation is made to permanently stabilise the species’ identification.

***Gymelloxes terea* (SCHAUS, 1892), comb. rev.**

Dalaca terea SCHAUS (1892: 330); Mexico, Paso de San Juan, SCHAUS leg.; coll. SCHAUS; [GP P. VIETTE 91513, type no. 18603]; [USNM]. — BERKKAU (1893: 190). — DRUCE (1898: 451; pl. 89, fig. 3 type [♂] dorsal). — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1294). — SCHRÖDER (1967: 340).

Gymelloxes terea: VIETTE (1952b: 28; fig. 2 ♂ [syn-]type gen.); coll. SCHAUS; GP P. VIETTE 2240.

Aepytyus (*Gymelloxes*) *terea*: NIELSEN & ROBINSON (1983: 19), syn.: *Aepytyus* (*Gymelloxes*) *muysca* (PFITZNER, 1914). — ROBINSON & NIELSEN (1984: 17), syn.: *Aepytyus* (*Gymelloxes*) *muysca* (PFITZNER, 1914).

Cibyra (*Gymelloxes*) *terea*: NIELSEN et al. (2000: 842), syn.: *Cibyra* (*Gymelloxes*) *muysca* (PFITZNER, 1914). — GREHAN (2010: 51; fig. d appendix).

***Gymelloxes trilinearis* (PFITZNER, 1914), comb. rev.**

Dalaca trilinearis PFITZNER (1914: 105); [Colombia], Soso-moco, 800 m, FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19). — SCHRÖDER (1967: 340); “holotype” [RECTE lectotype] ♂, SMFT 84; III. 1911; GP VIETTE no. 2097.

Dalaca trilinearides [sic]: PFITZNER (1937: 1294; pl. 99c [♂] dorsal).

Gymelloxes trilinearis: VIETTE (1952b: 28); lectotype.

[no genus] *trilinearides* [sic]: ZUKOWSKI (1954: 93).

Aepytyus (*Gymelloxes*) *trilinearis*: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Gymelloxes*) *trilinearis*: NIELSEN et al. (2000: 842).

***Hampsoniella* VIETTE, 1950, stat. rev.**

Type-species: *Dalaca assa* DRUCE, 1887 by original designation.

Aepytyus (*Hampsoniella*) VIETTE (1950a: 74); included species: *assa* (DRUCE, 1887), *equatorialis* VIETTE, 1950, *sladeni* (HAMPSON, 1903).

EDWARDS & HOPWOOD (1966: 116). — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Hampsoniella: VIETTE (1951b: 116). — VIETTE (1951d: 76). — PAULT (1953: 145); as synonym of *Aepytyus* HERRICH-SCHÄFFER, [1856]. — NYE & FLETCHER (1991: 141).

Cibyra (*Hampsoniella*): NIELSEN et al. (2000: 841). — GREHAN (2010: 49; fig. f appendix).

Remarks. *Hampsoniella* VIETTE, 1950 was described mainly based on the ♂ genitalia characters, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

***Hampsoniella assa* (DRUCE, 1887), comb. rev.**

Dalaca assa DRUCE (1887: 232; pl. 24, fig. 10 [♂] dorsal); Guatemala, Volcan de Atitlan, 2500-3000 feet, and Pantaleon, 1700 feet, CHAMPION [leg.]; BMNH. — KIRBY (1892: 886). — SCHAUS (1894: 236). — DRUCE (1898: 450). — WAGNER & PFITZNER (1911: 13). — DYAR (1915b: 350). — PFITZNER (1937: 1293; pl. 99c [♂] dorsal). — FORBES (1942: 406). — BIEZANKO (1961a: 8) [misidentification, unknown species].

Aepytyus (*Hampsoniella*) *assa*: VIETTE (1950a: 74; fig. 3 ♂ gen.) [error, unknown species; see VIETTE (1951c: 95)]. — NIELSEN

& ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Hampsoniella assa: VIETTE (1951c: 95; fig. 3 ([syn]type ♂ gen.)).

Cibyra (Hampsoniella) assa: NIELSEN et al. (2000: 841).

Hampsoniella equatorialis (VIETTE, 1950), **comb. rev.**

Aepytus (Hampsoniella) equatorialis VIETTE (1950a: 77; fig. 2 ♂ gen.); [holo-]type ♂, Ecuador, Bolivar, Balzapampa, M. DE MATHAN [leg.], IX. 1893–II. 1894; ex coll. OBERTHÜR, coll. BIEDERMANN; GP P. VIETTE no. 901; MNHN. — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Hampsoniella equatorialis: VIETTE (1951c: 95). — VIETTE (1951e: 1277).

Cibyra (Hampsoniella) equatorialis: NIELSEN et al. (2000: 841).

Hepialyxodes VIETTE, 1951, **stat. rev.**

Type-species: *Hepialyxodes rileyi* VIETTE, 1951, by original designation; monotypic.

VIETTE (1951e: 1278). — PACLT (1953: 145); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. — EDWARDS & HOPWOOD (1966: 120). — NYE & FLETCHER (1991: 146).

Aepytus (Hepialyxodes): NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Hepialyxodes): NIELSEN et al. (2000: 842). — GREHAN (2010: 44; fig. h appendix).

Cibyra (Hepialyxoides) [sic]; GREHAN (2010: 49).

Remarks. *Hepialyxodes* VIETTE, 1951 was described mainly based on the ♂ genitalia characters, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

Hepialyxodes rileyi VIETTE, 1951, **comb. rev.**

VIETTE (1951e: 1279; fig. 2 (♂ gen.)): holotype ♂, Brazil, São Paulo, Ypiranga [RECTE Ipiranga], III. 1932, R. SPITZ [leg.]; GP P. VIETTE no. 2300; BMNH.

Aepytus (Hepialyxodes) rileyi: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Hepialyxodes) rileyi: NIELSEN et al. (2000: 842).

Parapielus VIETTE, 1949

Type-species: *Pielus luteicornis* BERG, 1882, by original designation; monotypic.

VIETTE (1949a: 54); ♂ (1951d: 78). — PACLT (1953: 142). — DUMBLETON (1966: 971). — EDWARDS & HOPWOOD (1966: 213). — NIELSEN & ROBINSON (1983: 17, 100), syn.: *Lossbergiana* VIETTE, 1951. — MALLETT (1984: 77). — ROBINSON & NIELSEN (1984: 16), syn.: *Lossbergiana* VIETTE, 1951. — NIELSEN & SCOBLE (1986: 43). — NYE & FLETCHER (1991: 223). — NIELSEN et al. (2000: 840), syn.: *Lossbergiana* VIETTE, 1951. — SIMONSEN (2002: 65). — GREHAN (2010: 50).

= *Lossbergiana* VIETTE (1951a: 5); type-species: *Lossbergiana oberthuri* VIETTE, 1951, by original designation. — PACLT (1953: 143). — EDWARDS & HOPWOOD (1966: 153). — NIELSEN & ROBINSON (1983: 17, 100); as synonym of *Parapielus* VIETTE, 1949. — ROBINSON & NIELSEN (1984: 16); as synonym of *Parapielus* VIETTE, 1949. — NYE & FLETCHER (1991: 177). — NIELSEN et al. (2000: 840); as synonym of *Parapielus* VIETTE, 1949.

Parapielus heimlichii (URETA, 1956)

Hepialus heimlichii URETA (1956: 283); holotype ♂, Chile, Santiago, Aculeo, cerros bajos, 30. IV. 1955, HEIMLICH leg.; [GP ESN 2637, type no. 758]; coll. M.N. no. 6156; MNHC. — URETA (1957; pl. 1, fig. 7 ♀ dorsal). — CAMOUSSEIGHT (1980: 31).

Parapielus heimlichii: NIELSEN & ROBINSON (1983: 17, 105; figs. 68 (prelabium, labial palpus), 206–208 (♂ dorsal), 209 (♀ dorsal), 250 (♂ antennae flagellum), 331 (♂ holotype gen., ESN slide no. 2638), 362 (♀ gen.), 385 (bursa copulatrix), 413 (flight period), 430 (distribution)). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — DAPOTO et al. (2003: 100). — PASTRANA (2004: 6).

Parapielus luteicornis (BERG, 1882)

Pielus luteicornis BERG (1882: 218); [Chile], Patagonia (Fretum Magellanicum), Punta Arenas, Estrecho de Magallanes, 13. III. 1879; coll. STAUDINGER; [MACN]. — PFITZNER (1938: 1297; pl. 99h [♂] dorsal).

Epialus [sic] *luteicornis*: BERG (1882: 218).

Oxycanus luteicornis: KIRBY (1892: 893). — WAGNER & PFITZNER (1911: 21). — TINDALE (1981: 966).

Hepialus (Pielus) luteicornis: STAUDINGER (1899: 41; fig. 17 [♂]). — PAGENSTECHE (1902: 399).

Oxycanus niphadias (MEYRICK, 1890) [misidentification]: QUAIL (1900: 421); see TINDALE (1981: 966).

Pialus [sic] *luteicornis*: PAGENSTECHE (1902: 398).

Hepialus luteicornis: ENDERLEIN (1912: 90).

= *Pielus luteicornis* f. *popperi* PFITZNER (1938: 1297); [Chile and Argentina], Patagonia, Fuegia, Magallanes, Punta Arenas.

Parapielus luteicornis: VIETTE (1949a: 55; figs. 1 (labial palpus), 2–3 (antennal segment)). — ROBINSON (1977: 108). — NIELSEN & ROBINSON (1983: 17, 102; figs. 66 (prelabium, labial palpus), 69 (venation), 202 (♂ dorsal), 203 (♀ dorsal), 248 (♂ antennae flagellum), 274 (♂ foretibia), 275 (♀ foretibia), 329 (♂ gen.), 359 (♀ gen.), 383 (bursa copulatrix), 411 (flight period), 430 (distribution)); lectotype ♂, Chile, Magallanes, Punta Arenas, 13. III. 1879, BERG [leg.]; MACN, syn.: *Parapielus popperi* (PFITZNER, 1938). — ROBINSON & NIELSEN (1984: 16), syn.: *Parapielus popperi* (PFITZNER, 1938). — NIELSEN et al. (2000: 840), syn.: *Parapielus popperi* (PFITZNER, 1938). — PASTRANA (2004: 6). — GREHAN (2010: 44; fig. N appendix).

Parapielus popperi: NIELSEN & ROBINSON (1983: 17, 103); as synonym of *Parapielus luteicornis* (BERG, 1882). — ROBINSON & NIELSEN (1984: 16); as synonym of *Parapielus luteicornis* (BERG, 1882). — NIELSEN et al. (2000: 840); as synonym of *Parapielus luteicornis* (BERG, 1882).

Parapielus oberthuri (VIETTE, 1951)

Lossbergiana oberthuri VIETTE (1951a: 5; fig. 4 (♂ gen.)); [holo]type ♂, Chile, Valdivia, Arturo von LOSSBERG [leg.], 1904; ex coll. C. OBERTHÜR, coll. R. BIEDERMANN; GP P. VIETTE no. 1356; MNHN.

Parapielus oberthuri NIELSEN & ROBINSON (1983: 17, 104; figs. 67 (prelabium, labial palpus), 70 (venation), 204 (♂ dorsal), 205 (♀ dorsal), 249 (♂ antennae flagellum), 330 (♂ gen.), 360–361 (♀ gen.), 384 (bursa copulatrix), 412 (flight period), 430 (distribution)). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — PASTRANA (2004: 6).

Parapielus reedi (URETA, 1957)

Hepialus reedi URETA (1957: 163; fig. 4 ♂ gen.); holotype ♂, Chile, [Osorno, Lago Llanquihue], Puerto Octay, 23. II. 1956, E. OEHREN leg.; coll. M.N. no. 6200; MNHC. — CAMOUSSEIGHT (1980: 31).

Parapielus reedi: NIELSEN & ROBINSON (1983: 17, 107; figs. 210 (♂ holotype dorsal), 211 (♂ dorsal), 251 (♂ antennae flagellum), 276 (♂ foretibia), 332 (♂ holotype gen.), 414 (flight period), 431 (distribution)); GP ESN 2635, type no. 1054. — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840).

***Pfitzneriana* VIETTE, 1952**

Type-species: *Dalaca olivescens* PFITZNER, 1914, by original designation.

Pfitzneriana VIETTE (1952b: 29); included species: *olivescens* (PFITZNER, 1914), *vogli* VIETTE, 1952.

PACLT (1957: 51); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. — EDWARDS & HOPWOOD (1966: 223). — NIELSEN & ROBINSON (1983: 18), syn.: *Pseudophassus* WEYMER [ms.] (NEC PFITZNER, 1914). — ROBINSON & NIELSEN (1984: 17). — NYE & FLETCHER (1991: 232). — NIELSEN et al. (2000: 841). — GREHAN (2010: 43).

***Pfitzneriana allura* VIETTE, 1961**

VIETTE (1961: 5; fig. 6 ♂ gen.); holotype ♂, Bolivia, Santa Cruz, 500 m, 15. x. 1955, R. ZISCHKA leg.; GP P. VIETTE no. 3854; ZSBS. — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

***Pfitzneriana obliquestrigata* (STRAND, 1912), comb. n.**

Dalaca obliquestrigata STRAND (1912a: 156); [holo-]type ♂, Peru, oberer Madre de Dios; ZMHB. — STRAND (1914: 59, pl. IV, fig. 4 ♂ dorsal); ♀ (1927: 42). — PFITZNER (1937: 1294). — ZUKOWSKI (1954: 93). — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. The holotype was examined and this taxon is closely related to the type-species of *Pfitzneriana* VIETTE, 1952, *Dalaca olivescens* PFITZNER, 1914.

***Pfitzneriana olivescens* (PFITZNER, 1914)**

Dalaca olivescens PFITZNER (1914: 105); single specimen [= holotype], Colombia, Sosomoco, 800 m, FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19). — PFITZNER (1937: 1294; pl. 99e [♂] dorsal). — SCHRÖDER (1967: 340); holotype ♂, SMFT 85; i. 1911; GP P. VIETTE no. 2349.

Pfitzneriana olivescens: VIETTE (1952b: 30). — NIELSEN et al. (2000: 841), syn.: *Pfitzneriana boliviensis* VIETTE, 1961.

= *Pfitzneriana olivescens boliviensis* VIETTE (1961: 6); holotype ♂, Bolivia, [Cochabamba], Yungas del Palmar, 1000 m, 5. iii. 1949, R. ZISCHKA leg.; GP P. VIETTE no. 3549; ZSBS. — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Pfitzneriana olivescens olivescens: NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17).

Pfitzneriana boliviensis: NIELSEN et al. (2000: 841); as synonym of *Pfitzneriana olivescens* (PFITZNER, 1914).

***Pfitzneriana vogli* VIETTE, 1952**

VIETTE (1952b: 30; fig. 1 holotype gen.); holotype ♂, Venezuela, Caracas, Cerro Avila, 10.-15. iv. 1936, P. Cor. VOGL leg.; GP P. VIETTE no. 2396; ZSBS. — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

***Pfitzneriella* VIETTE, 1951**

Type-species: *Triodia remota* PFITZNER, 1906, by original designation; monotypic.

VIETTE (1951b: 116). — PACLT (1953: 142); ♀ (1957: 52). — EDWARDS & HOPWOOD (1966: 223). — SCHRÖDER (1967: 338). — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NYE & FLETCHER (1991: 232). — NIELSEN et al. (2000: 844). — GREHAN (2010: 47).

***Pfitzneriella lucicola* (MAASSEN, 1890)**

Triodia lucicola MAASSEN (1890: 137; pl. 4, fig. 16 [♂] dorsal); 4 specimens, Putzulagua near Latacunga, Ecuador, 3600 m; [ZMHB]. — BERTKAU (1891: 196). — PFITZNER (1906: 276); ♀

(1938: 1297; pl. 185f [♂] dorsal). — ZUKOWSKI (1954: 94).

Dalaca lucicola: KIRBY (1892: 886). — WAGNER & PFITZNER (1911: 14).

(no genus) *lucicola*: VIETTE (1951b: 116).

Pfitzneriella lucicola: NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

***Pfitzneriella monticola* (MAASSEN, 1890)**

Triodia monticola MAASSEN (1890: 136; pl. 4, fig. 14 [♂] dorsal); 11 specimens, Ecuador, Sincholagua, xi; [ZMHB]. — BERTKAU (1891: 196). — PFITZNER (1906: 276); ♀ (1938: 1297; pl. 185b [♂] dorsal). — ZUKOWSKI (1954: 94).

Dalaca monticola: KIRBY (1892: 886). — WAGNER & PFITZNER (1911: 14).

(no genus) *monticola*: VIETTE (1951b: 116).

Pfitzneriella monticola: NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844). — GREHAN (2010: 57; fig. A appendix).

***Pfitzneriella remota* (PFITZNER, 1906)**

Triodia remota PFITZNER (1906: 276); two specimens, Peru, Challabamba, 13400 feet, SCHULTZ leg.; coll. PFITZNER; [SMFL]; ♀ (1938: 1297; pl. 99e [♂] dorsal). — ZUKOWSKI (1954: 94).

Hepialus remotus [sic]; WAGNER & PFITZNER (1911: 10).

Pfitzneriella remota: VIETTE (1951b: 117; fig. 5 lectotype ♂ gen.). — SCHRÖDER (1967: 338); lectotype ♂, SMFT 1000; GP P. VIETTE no. 2100. — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

***Pfitzneriella similis* (ZUKOWSKI, 1954)**

Triodia similis ZUKOWSKI (1954: 94); [holo-]type, southern Peru, Rio Sondondo, 2400 m, 18. iv. 1936, Hamb. Südperu-Expedition; Hamburger Zoolog. Museum (specimen destroyed in war 1943).

Pfitzneriella similis: NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

***Phassus* WALKER, 1856**

Type-species: *Phassus argentiferus* WALKER, 1856, by subsequent designation by KIRBY (1892: 889).

Phassus WALKER (1856: 1566); included species: *argentiferus* WALKER, 1856, *giganteus* (HERRICH-SCHÄFFER, [1853]), *agrioides* WALKER, 1856, *transversus* WALKER, 1856, *tessellatus* (HERRICH-SCHÄFFER, [1854]), *signifer* WALKER, 1856.

DRUCE (1887: 232); ♀ (1892: 278). — HAMPSON ([1893]: 318); type-species: *Pharmacis huebneri* GEYER, ([1838]: pl. [53]) [incorrect designation]. — DRUCE (1898: 451). — QUAIL (1900: 422); [probably *Endoclita* C. & R. FELDER, 1874]. — WAGNER & PFITZNER (1911: 17). — WALSINGHAM (1915: 457). — STRAND (1916: 25). — LE CERF (1919: 469). — ANDRADE (1928: 451). — LIMA (1936: 282). — PFITZNER (1938: 1298). — NEAVE (1940: 701). — TINDALE (1941: 45). — PACLT (1944: 143). — BOURGOGNE (1949: 69), syn.: *Trichophassus* LE CERF, 1919. — VIETTE (1950c: 60); ♀ (1951d: 78). — PACLT (1953: 143). — SCHRÖDER (1967: 342). — SILVA et al. (1968: 202). — TINDALE (1981: 966). — NIELSEN & ROBINSON (1983: 18). — MALLET (1984: 76). — ROBINSON & NIELSEN (1984: 16). — GREHAN (1989: 805). — NYE & FLETCHER (1991: 234). — KRISTENSEN (1998: 62). — NIELSEN et al. (2000: 841). — SIMONSEN (2002: 65). — GREHAN & RAWLINS (2003: 733; fig. 1 (larva), figs. 2-8 (chaetotaxy)). — ARGUEDAS (2007: 48, 58). — CENGICANA (2008: 37). — GREHAN (2010: 46). — RAMOS-ELORDUY et al. (2011: 4).

Phasus [sic]: MONTE (1934: 211).

***Phassus aurigenus* PFITZNER, 1914**

PFITZNER (1914: 110); single specimen [= holotype], Costa Rica, Orosi, 1200 m; coll. PFITZNER; [SMFL]; ♂ (1938: 1299; pl. 99h [♂] dorsal). — VIETTE (1951e: 1282). — SCHRÖDER (1967: 342); holotype ♂, SMFT 47, GP VIETTE no. 2357. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus basirei* SCHAUS, 1890**

SCHAUS (1890: 46); two ♀♀, [Mexico], Coatepec; [GP P. VIETTE no. 91526, type no. 18788]; USNM. — KIRBY (1892: 890). — DRUCE (1898: 451). — WAGNER & PFITZNER (1911: 17). — PFITZNER (1938: 1300). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus championi* DRUCE, 1887**

DRUCE (1887: 233; pl. 24, fig. 11 [♀] dorsal); single specimen [= holotype], Guatemala, Purula, 4000 feet, CHAMPION leg.; BMNH. — KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 17). — PFITZNER (1938: 1300; pl. 185d ([♀] dorsal)). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus chrysodidyma* DYAR, 1915**

DYAR (1915a: 85); [holo-]type ♂ [RECTE ♀], Mexico, Zacualpan, vi. 1914, R. MÜLLER [leg.]; [GP VIETTE no. 91524], type no. 19334; USNM. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

***Phassus exclamationis* PFITZNER, 1938**

PFITZNER (1938: 1299); no data. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus huebneri* (GEYER, [1838])**

Pharmacis huebneri GEYER ([1838]: 53; pl. [53], figs. 1–2 [dorsal, ventral]).

= *Phassus argentiferus* WALKER (1856: 1566); Mexico; coll. HARTWEG; BMNH. — GERSTAECKER (1857: 425). — DRUCE (1887: 233); ♂ (1892: 279). — KIRBY (1892: 890). — DRUCE (1898: 451). — BARRETT (1900: 235). — SNODGRASS (1909: 565; figs. 149–152 (thorax), 202–203 (axilar sclerites)). — WAGNER & PFITZNER (1911: 17). — KÜNNETH (1914: 77). — WALSINGHAM (1915: 457); as synonym of *Phassus huebneri* (GEYER, [1838]). — PFITZNER (1938: 1300). — TINDALE (1941: 45; pl. 7, fig. 73 dorsal). — VIETTE (1950c: 61; fig. 4 ♂ gen.). — KRAUSS (1962: 135). — NIELSEN & ROBINSON (1983: 18); as synonym of *Phassus huebneri* (GEYER, [1838]). — ROBINSON & NIELSEN (1984: 16); as synonym of *Phassus huebneri* (GEYER, [1838]). — NIELSEN et al. (2000: 841); as synonym of *Phassus huebneri* (GEYER, [1838]). — DAY et al. (2003: 74).

Phassus huebneri: DRUCE (1887: 233). — HAMPSON ([1893]: 318). — WALSINGHAM (1915: 457), syn.: *Phassus argentiferus* WALKER, 1856. — SHEPARD (1930: 239, 241, 242, 243, 255; pl. 1, fig. 6 thorax). — NIELSEN & ROBINSON (1983: 18), syn.: *Phassus argentiferus* WALKER, 1856, *Phassus pedipogon* STRAND, 1916. — ROBINSON & NIELSEN (1984: 16), syn.: *Phassus argentiferus* WALKER, 1856, *Phassus pedipogon* STRAND, 1916. — NIELSEN et al. (2000: 841), syn.: *Phassus argentiferus* WALKER, 1856, *Phassus pedipogon* STRAND, 1916.

= *Phassus pedipogon* STRAND (1916: 25; pl. 15, fig. 6 [♀] dorsal); single specimen [= holotype ♀], Costa Rica; coll. NIEPELT; [BMNH]. — STRAND (1927: 42). — PFITZNER (1938: 1299). — NIELSEN & ROBINSON (1983: 18); as synonym of *Phassus huebneri* (GEYER, [1838]). — ROBINSON & NIELSEN (1984: 16); as synonym of *Phassus huebneri* (GEYER, [1838]). — NIELSEN et al. (2000: 841); as synonym of *Phassus huebneri* (GEYER, [1838]).

Phassus triangularis f. *huebneri*: PFITZNER (1938: 1299; pl. 100c dorsal).

***Phassus marcius* DRUCE, 1892**

DRUCE (1892: 278); Mexico, near Durango City, BECKER [leg.]; [BMNH]. — KIRBY (1892: 890). — BERTKAU (1893: 190). — DRUCE (1898: 451; pl. 89, fig. 4 dorsal). — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1300; pl. 100a dorsal). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus n-signatus* WEYMER, 1907**

WEYMER (1907: 37); single ♀ [= holotype], Guatemala; coll. WERNICKE. — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1300). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Phassus triangularis [misidentification]: PFITZNER (1938; pl. 100a [♀] dorsal).

***Phassus phalerus* DRUCE, 1887**

DRUCE (1887: 233; pl. 24, fig. 8 ♂ dorsal); single ♂ [= holotype], Mexico, [Vera Cruz], Jalapa, HöGE [leg.]; [BMNH]. — KIRBY (1892: 890). — DRUCE (1898: 451). — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1300; pl. 185f [♂] dorsal). — NIELSEN & ROBINSON (1983: 18). — MALLETT (1984: 77). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841). — CENGICANA (2007: 57).

***Phassus pharus* (DRUCE, 1887)**

Hepialus pharus DRUCE (1887: 232; pl. 24, fig. 12 ♂ dorsal); Guatemala, Las Mercedes, 3000 feet, and Dueñas, CHAMPION [leg.]; Costa Rica, Irazu, 6000–7000 feet, ROGERS [leg.]; [BMNH]. — KIRBY (1892: 884). — WAGNER & PFITZNER (1911: 9). — WILLIAMS (1935: 292; figs. 1.1 (dorsal), 1.2 (egg), 1.3 (larva), 1.4–5 (pupa), 1.6 (resting), 1.7 (♂ hindleg), 2–3 (behaviour)).

Hepialus (*Hepialus*) *pharus*: PFITZNER (1937: 1291).

Phassus pharus: NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus rosulentus* WEYMER, 1907**

WEYMER (1907: 35); single ♂ [= holotype], Mexico, [Vera Cruz], Jalapa; coll. FICKE. — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1299). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

***Phassus triangularis* EDWARDS, 1885**

EDWARDS (1885: 129); Mexico, [Vera Cruz], Jalapa, W. SCHAUS leg. — DRUCE (1887: 233). — SCHAUS (1888: 64). — KIRBY (1892: 890). — PACKARD (1895: 74; fig. 34 pupa). — DRUCE (1898: 451; pl. 89, fig. 1 [♀] dorsal). — BARRETT (1900: 235). — HEADLEE (1907: 285; pl. 60, fig. 9 hindwing [error]). — SNODGRASS (1909: 565; figs. 153–154 thorax). — WAGNER & PFITZNER (1911: 19). — DYAR (1917: 132). — PFITZNER (1938: 1299). — NIELSEN & ROBINSON (1983: 18), syn.: *Phassus triangularides* (PFITZNER, 1938). — ROBINSON & NIELSEN (1984: 16), syn.: *Phassus triangularides* (PFITZNER, 1938). — HILJE et al. (1992a: 152). — NIELSEN et al. (2000: 841), syn.: *Phassus triangularides* (PFITZNER, 1938). — ARGUEDAS & ESPINOZA (2007: 3); fig. 7 (larva). — ARGUEDAS (2007: 5, 57). — GREHAN (2010: 62; fig. U, appendix). — RAMOS-ELORDUY et al. (2011: 4).

= *Phassus triangularis* f. *triangularides* PFITZNER (1938: 1299; pl. 100c [♀] dorsal); Mexico.

Phassus triangularides: NIELSEN & ROBINSON (1983: 18); as synonym of *Phassus triangularis* EDWARDS, 1885. — ROBINSON & NIELSEN (1984: 16); as synonym of *Phassus triangularis* EDWARDS, 1885. — NIELSEN et al. (2000: 841); as synonym of *Phassus triangularis* EDWARDS, 1885.

***Phialuse* VIETTE, 1961**

Type-species: *Phialuse palmar* VIETTE, 1961, by original designation; monotypic.

VIETTE (1961: 4). — EDWARDS & VEVERS (1975: 258). — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NYE & FLETCHER (1991: 235). — NIELSEN et al. (2000: 843). — GREHAN (2010: 50).

***Phialuse palmar* VIETTE, 1961**

VIETTE (1961: 4; fig. 5 ♂ gen.); holotype ♂, Bolivia, [Cochabamba], Yungas del Palmar, 1500 m, 22. VII. 1952, R. ZISCHKA leg.; GP P. VIETTE no. 3785; ZSBS. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 843).

***Philoenia* KIRBY, 1892, stat. rev.**

Type-species: *Pharmacis lagopus* MÖSCHLER, 1877, by original designation; monotypic.

KIRBY (1892: 885). — NEAVE (1940: 711). — PACLT (1944: 143); ♂ (1953: 145); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. — NYE & FLETCHER (1991: 236).

Philaenia [sic]: WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1293). — VIETTE (1951b: 116); ♂ (1951d: 78). — SCHRÖDER (1967: 340). — NYE & FLETCHER (1991: 235).

Aepytus (Philaenia) [sic]: VIETTE (1951a: 1). — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Philoenia): NIELSEN et al. (2000: 843). — GREHAN (2010: 49).

Remarks. *Philoenia* KIRBY, 1892 was characterised by VIETTE (1951b) mainly based on the ♂ genitalia, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

***Philoenia brasiliensis* VIETTE, 1952, comb. rev.**

Philaenia [sic] *brasiliensis* VIETTE (1952a: 143; fig. 6 saccus); holotype ♂, Brazil, [Rio de Janeiro], Petrópolis, 31. III. 1913, J. G. FOETTERLE leg.; GP P. VIETTE no. 2259; NHMW.

Aepytus (Philaenia) [sic] *brasiliensis*: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Philoenia) brasiliensis: NIELSEN et al. (2000: 843).

***Philoenia fasslii* (PFITZNER, 1914), comb. rev.**

Dalaca fasslii PFITZNER (1914: 106); 2 ♂♂, 1 ♀, Colombia, Sosomoco, 800 m, [III. 1910], and Villavicencio, 450 m, [IV. 1910], FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19). — PFITZNER (1937: 1296; pl. 99f [♂, ♀] dorsal).

Philaenia [sic] *fasslii*: VIETTE (1951b: 118; fig. 4 lectotype ♂ gen.); lectotype ♂, gen. prep. VIETTE nr. 1395; ♂ (1951e: 1276); ♂ (1952a: 143). — SCHRÖDER (1967: 340); lectotype ♂, SMFT 93; III. 1910.

Aepytus (Philaenia) [sic] *fasslii*: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Philoenia) fasslii: NIELSEN et al. (2000: 843).

***Philoenia guyanensis* (VIETTE, 1951), comb. rev.**

Aepytus (Philaenia) [sic] *guyanensis* VIETTE (1951a: 3); [holo-]type ♂, French Guyana; coll. C. BAR, in coll. C. OBERTHÜR, coll. R. BIEDERMANN; GP P. VIETTE no. 1346; MNHN. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Philaenia [sic] *guyanensis*: VIETTE (1952a: 143).

Cibyra (Philoenia) guyanensis: NIELSEN et al. (2000: 843).

***Philoenia indicata* (STRAND, 1912), comb. n.**

Dalaca indicata STRAND (1912b: 100); [holo-]type ♂, Ecuador, Macas, [1905-06]; coll. W. NIEPELT; [GP P. VIETTE no.

2025]; [BMNH]. — STRAND (1914: 59; pl. 11, fig. 13 ♂ dorsal). — STRAND (1927: 42). — PFITZNER (1937: 1294). — ZUKOWSKI (1954: 93).

Aepytus (Philaenia) [sic] *indicata*: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Philoenia) indicata: NIELSEN et al. (2000: 843).

***Philoenia lagopus* (MÖSCHLER, 1877), comb. rev.**

Pharmacis lagopus MÖSCHLER (1877: 670; pl. 9, fig. 34 [♂] dorsal); 2 ♂♂, Inner Surinam; [ZMHB].

Philoenia lagopus: KIRBY (1892: 885).

Philaenia [sic] *lagopus*: WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1293; pl. 99c [♂] dorsal). — VIETTE (1952a: 143). — BIEZANKO (1961a: 8 [misidentification, unknown species]).

Aepytus (Philaenia) [sic] *lagopus*: VIETTE (1951a: 3; fig. 1 ♂ gen.). — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Philoenia) lagopus: NIELSEN et al. (2000: 843).

***Philoenia saguanmachica* (PFITZNER, 1914), comb. rev.**

Dalaca saguanmachica PFITZNER (1914: 110); East Colombia, Buenavista, 1200 m, FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19). — PFITZNER (1937: 1296; pl. 99g [♂] dorsal).

Philaenia [sic] *saguanmachica*: VIETTE (1951b: 118; fig. 6 ♂ gen. [lectotype]); lectotype ♂; GP P. VIETTE no. 2101; ♂ (1952a: 143). — SCHRÖDER (1967: 341); lectotype ♂, SMFT 1011.

Aepytus (Philaenia) [sic] *saguanmachica*: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Philoenia) saguanmachica: NIELSEN et al. (2000: 843).

***Philoenia thisbe* (DRUCE, 1901), comb. n.**

Dalaca thisbe DRUCE (1901: 437); Colombia, Don Amo; coll. DRUCE; [BMNH]. — PFITZNER (1937: 1296).

= *Dalaca thisbe* f. *hemichrysea* PFITZNER (1937: 1296; pl. 99g [♂] dorsal); East Colombia, Sosomoco, 800 m, FASSL [leg.]; [SMFL].

Dalaca hemichrysea: SCHRÖDER (1967: 339); III. 1911, FASSL leg.; "holotype" [RECTE lectotype] ♂, SMFT 91, GP VIETTE no. 2352.

Aepytus (Philaenia) [sic] *thisbe*: NIELSEN & ROBINSON (1983: 20), syn.: *Aepytus (Philaenia)* [sic] *hemichrysea* (PFITZNER, 1937). — ROBINSON & NIELSEN (1984: 17), syn.: *Aepytus (Philaenia)* [sic] *hemichrysea* (PFITZNER, 1937).

Aepytus (Philaenia) [sic] *hemichrysea*: NIELSEN & ROBINSON (1983: 20); as synonym of *Aepytus (Philaenia)* [sic] *thisbe* (DRUCE, 1901). — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytus (Philaenia)* [sic] *thisbe* (DRUCE, 1901).

Cibyra (Philoenia) thisbe: NIELSEN et al. (2000: 843), syn.: *Cibyra (Philoenia) hemichrysea* (PFITZNER, 1937).

Cibyra (Philoenia) hemichrysea: NIELSEN et al. (2000: 843); as synonym of *Cibyra (Philoenia) thisbe* (DRUCE, 1901).

***Pseudodalaca* VIETTE, 1950, stat. rev.**

Type-species: *Dalaca sarta* SCHAUS, 1894, by original designation.

Aepytus (Pseudodalaca) VIETTE (1950a: 74); included species: *sarta* (SCHAUS, 1894), *gugelmanni* VIETTE, 1950.

NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17).

Pseudodalaca: VIETTE (1951c: 95); type-species *Pseudodalaca gugelmanni* VIETTE, 1950, by subsequent designation [incorrect designation]. — VIETTE (1951b: 116); ♂ (1951d: 78). —

PACLT (1953: 145); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. — EDWARDS & HOPWOOD (1966: 244). — NYE & FLETCHER (1991: 257).

Cibyra (*Pseudodalaca*): NIELSEN et al. (2000: 842). — GREHAN (2010: 49).

Remarks. *Pseudodalaca* VIETTE, 1950 was described by VIETTE (1950a) mainly based on the ♂ genitalia characters, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

***Pseudodalaca gugelmanni* (VIETTE, 1950), comb. rev.**

Aepytus (*Pseudodalaca*) *gugelmanni* VIETTE (1950a: 78; fig. 6 ♂ gen.); [holo-]type ♂, Mexico, [Vera Cruz], Misantla, W. GÜGELMANN [leg.], iv–v. 1912; ex coll. OBERTHÜR, coll. R. BIEDERMANN; GP P. VIETTE no. 918; MNHN. — NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Pseudodalaca gugelmanni: VIETTE (1951c: 95); ♂ (1951e: 1277).

Cibyra (*Pseudodalaca*) *gugelmanni*: NIELSEN et al. (2000: 842). — GREHAN (2010: 53; fig. j appendix).

***Pseudodalaca mexicanensis* VIETTE, 1953, comb. rev.**

VIETTE (1953a: 20; fig. 1 holotype ♂ gen.); holotype ♂, [Mexico], [Vera Cruz], Jalapa; coll. C. OBERTHÜR, coll. R. BIEDERMANN; GP P. VIETTE no. 920; MNHN; ♂ (1950a: 78; fig. 5 ♂ gen., as *Aepytus* (*Pseudodalaca*) *serta* (SCHAUS, 1894)).

Aepytus (*Pseudodalaca*) *mexicanensis*: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Pseudodalaca*) *mexicanensis*: NIELSEN et al. (2000: 842).

***Pseudodalaca sertae* (SCHAUS, 1894), comb. n.**

Dalaca sertae SCHAUS (1894: 236); ♂, Mexico, [Vera Cruz], Jalapa; [GP P. VIETTE 91514, type no. 18601]; USNM. — DRUCE (1898: 450; pl. 89, fig. 2 [syn-]type ♂ dorsal). — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1294).

Aepytus (*Pseudodalaca*) *sertae*: VIETTE (1950a: 78; fig. 5 ♂ gen. [error, *P. mexicanensis* VIETTE, 1953]). — ROJAS & CHACÓN (1980: 63). — AGUIAR-MENEZES et al. (2002: 377). — VERGARA (2005: 98; figs. 15 (biology), 16 (♂ dorsal)).

Hampsoniella sertae: VIETTE (1951c: 95; fig. 1 [syn-]type ♂ gen.).

Aepytus (*Hampsoniella*) *sertae*: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Hampsoniella*) *sertae*: NIELSEN et al. (2000: 841).

Remarks. *Dalaca sertae* SCHAUS, 1894 was designated the type-species of *Pseudodalaca* VIETTE, 1950 by original designation.

***Pseudophassus* PFITZNER, 1914, stat. rev.**

Type-species: *Pseudophassus mahagoniatus* PFITZNER, 1914, by subsequent designation by VIETTE (1961: 6).

Pseudophassus PFITZNER (1914: 110); included species: *songoensis* PFITZNER, 1914, *mahagoniatus* PFITZNER, 1914.

VIETTE (1961: 6), syn.: *Parana* VIETTE, 1951. — SCHRÖDER (1967: 342). — NYE & FLETCHER (1991: 258).

Pseudophassus WEYMER [ms.]: PFITZNER (1938: 1301). — VIETTE (1950d: 165); type-species: *Hepialus prosopus* DRUCE, 1901, by subsequent designation [incorrect designation]. — PACLT (1953: 145), syn.: *Druceiella* VIETTE, 1949. — ZUKOWSKI (1954: 94). — EDWARDS & HOPWOOD (1966: 247). — NIELSEN & ROBINSON (1983: 18); as synonym of *Pfitzneriana* VIETTE, 1952; *praeocc.*: PFITZNER, 1914 [Hepialidae].

Phassus (*Pseudophassus*): PACLT (1944: 143).
= *Aepytus* (*Parana*) VIETTE (1950a: 75); type-species: *Aepytus* (*Parana*) *philiponi* VIETTE, 1950, by original designa-

tion. — NIELSEN & ROBINSON (1983: 20); as synonym of *Aepytus* (*Tricladia*) C. & R. FELDER, 1874. — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytus* (*Tricladia*) C. & R. FELDER, 1874.

Parana: VIETTE (1951b: 116). — PACLT (1953: 145); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. — VIETTE (1961: 6); as synonym of *Pseudophassus* PFITZNER, 1914. — EDWARDS & HOPWOOD (1966: 211). — NYE & FLETCHER (1991: 223). — *Præocc.*: *Parana* NIXON, 1943 [Hymenoptera]; ♂ (1951d: 78).

Aepytus (*Pseudophassus*): NIELSEN & ROBINSON (1983: 18, 20); as synonym of *Aepytus* (*Tricladia*) C. & R. FELDER, 1874. — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytus* (*Tricladia*) C. & R. FELDER, 1874.

Cibyra (*Pseudophassus*): NIELSEN et al. (2000: 843); as synonym of *Cibyra* (*Tricladia*) C. & R. FELDER, 1874.

Cibyra (*Parana*): NIELSEN et al. (2000: 843); as synonym of *Cibyra* (*Tricladia*) C. & R. FELDER, 1874.

Remarks. The type-species of *Pseudophassus* PFITZNER, 1914 was examined, and based on the ♂ genitalia it is clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

***Pseudophassus mahagoniatus* PFITZNER, 1914, comb. rev.**

PFITZNER (1914: 110); [Bolivia], Rio Songo, 1912, FASSL [leg.]; coll. PFITZNER; [SMFL]; ♂ (1938: 1301; pl. 99h [♂] dorsal). — SCHRÖDER (1967: 342); lectotype ♂, SMFT 49; GP VIETTE no. 2356.

Parana mahagoniatus: VIETTE (1951e: 1277).

Aepytus (*Tricladia*) *mahagoniatus*: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Tricladia*) *mahagoniatus*: NIELSEN et al. (2000: 843). — GREHAN (2010: 54; fig. m (appendix)).

***Pseudophassus philiponi* (VIETTE, 1950), comb. n.**

Aepytus (*Parana*) *philiponi* VIETTE (1950a: 80; fig. 7 ♂ gen.); [holo-]type ♂, Brazil, Para [RECTE Pará], 1927, Comte R. PHILIPON [leg.]; GP P. VIETTE no. 1360; MNHN.

Parana philiponi: VIETTE (1951e: 1277).

Aepytus (*Tricladia*) *philiponi*: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Tricladia*) *philiponi*: NIELSEN et al. (2000: 843).

Remarks. Based on the holotype examination, *Aepytus* (*Parana*) *philiponi* VIETTE is closely related to *Pseudophassus mahagoniatus* PFITZNER, 1914, the type-species of *Pseudophassus* PFITZNER, 1914.

***Pseudophilaenia* VIETTE, 1951, stat. rev.**

Type-species: *Philaenia* [sic] *lagopus* f. *omagua* PFITZNER, 1937, by original designation; monotypic.

VIETTE (1951b: 116). — PACLT (1953: 145); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1856]. — EDWARDS & HOPWOOD (1966: 247). — SCHRÖDER (1967: 341). — NYE & FLETCHER (1991: 258).

Aepytus (*Pseudophilaenia*): NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (*Pseudophilaenia*): NIELSEN et al. (2000: 843). — GREHAN (2010: 44).

Remarks. *Pseudophilaenia* VIETTE, 1951 was described mainly based on the ♂ genitalia characters, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

***Pseudophilaenia omagua* (PFITZNER, 1937), comb. rev.**

Philaenia [sic] *lagopus* f. *omagua* PFITZNER (1937: 1293); [Brazil, Amazonas], Upper Rio Negro and [Peru, Loreto], Amazons (Pebas); [SMFL].

Pseudophilaenia omagua: VIETTE (1951b: 117; figs. [♂] lecto-

type]: 1 (8th tergite), 2 (gen.), 3 (8th sternite)); lectotype ♂, GP VIETTE nr. 1393; coll. PFITZNER; ♀ (1951e: 1276). — SCHRÖDER (1967: 341); lectotype ♂, SMFT 1029.

Aepytus (Pseudophilaenia) omagua: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Pseudophilaenia) omagua: NIELSEN et al. (2000: 843). — GREHAN (2010: 53; fig. i appendix).

Puermytrans VIETTE, 1951

Type-species: *Puermytrans chiliensis* VIETTE, 1951, by original designation; monotypic.

VIETTE (1951e: 1273). — PAULT (1953: 143); ♀ (1957: 51). — EDWARDS & HOPWOOD (1966: 251). — NIELSEN & ROBINSON (1983: 17, 95). — ROBINSON & NIELSEN (1984: 16). — NYE & FLETCHER (1991: 263). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN & RAWLINS (2003: 734). — GREHAN (2010: 50).

Puermytrans chiliensis VIETTE, 1951

VIETTE (1951e: 1274; fig. 1 ♂ gen.); holotype ♂, Chile, WALKER; coll. H. DRUCE; GP P. VIETTE 2294; BMNH. — NIELSEN & ROBINSON (1983: 17, 98; figs. 58 (prelabium, labial palpus), 59 (venation), 60–63 (wing scent organ), 64–65 (hind leg, abdomen), 198–199 (♂ dorsal), 200–201 (♀ dorsal), 246 (♂ antennae flagellum), 247 (♀ antennae flagellum), 273 (♂ foretibia), 327–328 (♂ gen.), 357–358 (♀ gen.), 382 (bursa copulatrix), 410 (flight period), 429 (distribution)). — MALLET (1984: 77). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN (2010: 63; fig. Z appendix).

Roseala VIETTE, 1950

Type-species: *Roseala bourgognei* VIETTE, 1950, by original designation; monotypic.

VIETTE (1950c: 53); ♀ (1951d: 79). — PAULT (1953: 145), syn.: *Thiastyx* VIETTE, 1951, *Xytraps* VIETTE, 1951 [see under *Cibyra* WALKER, 1856]. — EDWARDS & HOPWOOD (1966: 261). — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NYE & FLETCHER (1991: 269). — NIELSEN et al. (2000: 843).

= *Thiastyx* VIETTE (1951e: 1275); type-species: *Thiastyx catharinae* VIETTE, 1951 by original designation. — PAULT (1953: 145); as synonym of *Roseala* VIETTE, 1950. — EDWARDS & HOPWOOD (1966: 293). — NYE & FLETCHER (1991: 303).

Aepytus (Thiastyx): NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Thiastyx): NIELSEN et al. (2000: 842). — GREHAN (2010: 49); as a synonym of *Cibyra (Roseala)* VIETTE, 1950.

Cibyra (Roseala): GREHAN (2010: 50), syn.: *Cibyra (Thiastyx)* VIETTE, 1951

Roseala tessellatus (HERRICH-SCHÄFFER, [1854]), **comb. n.**

Epialus [sic] *tessellatus* HERRICH-SCHÄFFER ([1854] (BOISDUVAL *in litt.*); cover, [pl. 31], fig. 147 [♀] dorsal); Nov. Holl.; [MNHN]; ♀ ([ix. 1856]: 5; fig. 147; ♀ ([1858]: 57, 79; fig. 147; Port Natal [Durban, South Africa – error]).

= *Phassus agrionides* WALKER (1856: 1567); Brazil; coll. BECKER; BMNH. — KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 17). — PFITZNER (1938: 1299). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841); **syn. n.**

Phassus tessellatus: WALKER ([v.] 1856: 1568). — KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 19). — JANSE (1917: 219). — LE CERF (1919: 470). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16).

= *Roseala bourgognei* VIETTE (1950c: 54; fig. 3 ♂ gen.); [ho-

lo-]type ♂, Brazil, Petrópolis, 18. iv. [19]07, J. G. FOETTERLE [leg.]; GP P. VIETTE no. 1363; NHMW. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 843); **syn. n.**

= *Thiastyx catharinae* VIETTE (1951e: 1276; fig. 3 ♂ gen.); holotype ♂, Brazil, Santa Catarina, Nova Bremen [Dalbergia], Rio Laeiss [RECTE Rio Lais], iv. 1936, F. H. HOFFMANN [leg.]; GP P. VIETTE no. 2370; ex coll. Tring Museum; BMNH; **syn. n.**

Aepytus (Thiastyx) catharinae: NIELSEN & ROBINSON (1983: 19, 44). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Thiastyx) catharinae: NIELSEN et al. (2000: 842).

Phassus tessellatus [sic]: NIELSEN et al. (2000: 841).

Cibyra (Roseala) catharinae: GREHAN (2010: 53; fig. k appendix).

Remarks. One syntype ♀ in the BMNH of *Phassus agrionides* WALKER, 1856 that bears a label “holotype” is here designated lectotype; it has the following labels: “holotype/ 3. *Phassus agrionides*/ 46 46/ syn. of *tessellatus* HS. new synonymy 1936 N. B. TINDALE in MS/ ... Bras 1. One syntype ♀ in the MNHN of *Epialus* [sic] *tessellatus* HERRICH-SCHÄFFER, [1854] that bears a label “type” is here designated lectotype; it has the following labels: / Type/ *Epialus tessellatus* HS. Samml. aussereurop. Schmett. 1853, pl. 31. fig. 147/ *Hepialus tessellatus* H.S. ♀ Type/ ex musaeo D. BOISDUVAL/ coll. R. BIEDERMANN/ ex-collection OBERTHUR, R. BIEDERMANN det., Muséum Paris/ Photo, det. E. S. NIELSEN 1984. — The types of *Epialus* [sic] *tessellatus* HERRICH-SCHÄFFER, [1854], *Phassus agrionides* WALKER, 1856, *Roseala bourgognei* VIETTE, 1950, and *Thiastyx catharinae* VIETTE, 1951 were examined and through examination of large series, morphology and mtDNA barcode sequences, all are synonymies. The two designations are made to permanently stabilise the species’ identification.

Schaefferiana VIETTE, 1950, **stat. rev.**

Type-species: *Epialus* [sic] *epigramma* HERRICH-SCHÄFFER, [1854], by original designation.

Schaefferiana VIETTE (1950c: 58); included species: *epigramma* (HERRICH-SCHÄFFER, [1854]), *jeanneli* VIETTE, 1950, *biedermanni* VIETTE, 1950.

VIETTE (1951d: 79). — PAULT (1953: 143). — EDWARDS & HOPWOOD (1966: 266). — NYE & FLETCHER (1991: 273).

Aepytus (Schaefferiana): NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Schaefferiana): NIELSEN et al. (2000: 842). — GREHAN (2010: 44).

Remarks. *Schaefferiana* VIETTE, 1950 was described mainly based on the ♂ genitalia characters, clearly distinct from the other genera. Therefore, we re-instate it here as a separate genus.

Schaefferiana epigramma (HERRICH-SCHÄFFER, [1854]), **comb. rev.**

Epialus [sic] *epigramma* HERRICH-SCHÄFFER ([1854] (BOISDUVAL *in litt.*); cover; pl. [31], fig. 146 [♂] dorsal); [Brazil]; [MNHN]; ♀ ([1858]: 79).

Dalaca epigramma: WALKER ([v.] 1856: 1561). — GERSTAECKER (1857: 425). — PFITZNER (1937: 1293; pl. 185a [♂] dorsal). — BIEZANKO et al. (1957: 9). — BIEZANKO (1961a: 8); ♀ (1961b: 8).

Triodia epigramma: HERRICH-SCHÄFFER ([ix. 1856]: 5; fig. 141 [RECTE 146]); ♀ ([1858]: 56); Brazil.

Aepytus epigramma: KIRBY (1892: 887).

Hepialus epigramma: WAGNER & PFITZNER (1911: 4).

Schaefferiana epigramma: VIETTE (1950c: 58; fig. 9 ♂ gen.); ♀ (1951c: 96).

Schaefferiana epigramme [sic]: VIETTE (1950c: 58).

Aepytus (Schaefferiana) epigramma: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Schaefferiana) epigramma: NIELSEN et al. (2000: 842). — GREHAN (2010: 53; fig. 1 appendix).

Schaefferiana simplex VIETTE, 1956, **comb. rev.**

VIETTE (1956: 378; fig. 5 ♂ gen.); holotype ♂, Brazil, Minas Gerais, San Jacintho Valley, Teophilo Ottoni [RECTE Teófilo Otoni], 1907–8, R. BIRCH [leg.]; GP P. VIETTE no. 2991; BMNH.

Aepytus (Schaefferiana) simplex: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Schaefferiana) simplex: NIELSEN et al. (2000: 842).

Schausiana VIETTE, 1950

Type-species: *Phassus trojesa* SCHAUS, 1901, by original designation; monotypic.

VIETTE (1950a: 80); ♂ (1951d: 79). — PACLT (1953: 143). — EDWARDS & HOPWOOD (1966: 266). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NYE & FLETCHER (1991: 273). — NIELSEN et al. (2000: 841). — GREHAN (2010: 47; fig. V appendix).

Schausiana trojesa (SCHAUS, 1901)

Phassus trojesa SCHAUS (1901: 76); Mexico, Trojes; [type no. 18613]; [USNM]. — WAGNER & PFITZNER (1911: 19). — PFITZNER (1938: 1300; pl. 100d dorsal). — VIETTE (1950b: 190).

Schausiana trojesa: VIETTE (1950a: 80; fig. 8 ♂ gen.). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

Phassus trajesa [sic]: RAMOS-ELORDUY et al. (2011: 4).

Trichophassus LE CERF, 1919

Type-species: *Epiolus* [sic] *giganteus* HERRICH-SCHÄFFER, [1853], by original designation; monotypic.

LE CERF (1919: 470). — NEAVE (1940: 545). — BOURGOGNE (1949: 69); as synonym of *Phassus* WALKER, 1856. — VIETTE (1949b: 72); ♂ (1951d: 79). — PACLT (1953: 142). — DUMBLETON (1966: 926, 927, 940, 971). — NIELSEN & ROBINSON (1983: 18). — MALLETT (1984: 77). — ROBINSON & NIELSEN (1984: 16). — GREHAN (1989: 805). — NYE & FLETCHER (1991: 313). — NIELSEN et al. (2000: 840). — SIMONSEN (2002: 65). — GREHAN & RAWLINS (2003: 734). — GREHAN (2010: 43).

Trichophassus [sic]: GREHAN (1984: 52).

Trichophassus giganteus (HERRICH-SCHÄFFER, [1853])

Epiolus [sic] *giganteus* HERRICH-SCHÄFFER ([1853] (BOISDUVAL, *in litt.*): cover; pl. 10, fig. 45 [♀] dorsal); “Amer. aequin.”; ♂ ([1858]: 78).

Phassus giganteus: WALKER (1856: 1567). — GERSTAECKER (1857: 425). — KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 18). — HOFFMANN (1931: 3). — PFITZNER (1938: 1301; pl. 100d [♀] dorsal). — LIMA (1945: 12, 146; figs. 7 thorax, 45 [♀] dorsal, 46 venation). — OITICICA FILHO (1947: 389; figs. 7–12 ♀ gen.). — BRIQUELOT (1956: 1; figs. 1 (larva), 2 (pupae), 3 (♂ dorsal), 4 (♂ resting), 5 (♀ dorsal), 6a, b (scales), 7 (venation), 8 (jugum), 9 (♀ gen.), 13–18, 20 (biology)). — BIEZANKO (1961: 8). — SILVA et al. (1968: 203). — PASTRANA (2004: 6).

Epiolus [sic] *giganteus*: HERRICH-SCHÄFFER ([1856]: 5; fig. 45); ♂ ([1858]: 57).

= *Phassus hayeki* FOETTERLE (1903: 649; fig. 1 ♂ dorsal); 3 ♂♂, [Brazil, Rio de Janeiro], Petrópolis. — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1301; pl. 185b [♂] dorsal).

Trichophassus giganteus: LE CERF (1919: 470). — VIETTE (1949b: 72; figs. 1–2 (antenna), 3 (♂ gen.)), syn.: *Trichophas-*

sus hayeki (FOETTERLE, 1903). — BOURGOGNE (1949: 69, 76; figs. 11–12 ♀ gen.). — DUMBLETON (1966: 925). — NIELSEN & ROBINSON (1983: 18), syn.: *Trichophassus hayeki* (FOETTERLE, 1903). — ROBINSON & NIELSEN (1984: 16), syn.: *Trichophassus hayeki* (FOETTERLE, 1903). — SBORDONI & FORESTIERO (1985: 88). — BUZZI (1994: 29, 141, 204). — DUGDALE (1994: 12). — KRISTENSEN (1998: 62). — NIELSEN et al. (2000: 840), syn.: *Trichophassus hayeki* (FOETTERLE, 1903). — SIMONSEN (2002: 65). — BUZZI (2009: 71, 298, 484). — GREHAN (2010: 45).

Trichophassus hayeki: VIETTE (1949b: 72); as synonym of *Trichophassus giganteus* (HERRICH-SCHÄFFER, [1853]). — NIELSEN & ROBINSON (1983: 18); as synonym of *Trichophassus giganteus* (HERRICH-SCHÄFFER, [1853]). — ROBINSON & NIELSEN (1984: 16); as synonym of *Trichophassus giganteus* (HERRICH-SCHÄFFER, [1853]). — NIELSEN et al. (2000: 840); as synonym of *Trichophassus giganteus* (HERRICH-SCHÄFFER, [1853]).

Tricladia C. & R. FELDER, 1874, **stat. rev.**

Type-species: *Tricladia umbrifera* C. & R. FELDER, 1874, by monotypy.

(*Tricladia* C. & R. FELDER, 1874 is senior homonym of *Tricladia* OBERTHÜR, 1894 [Lepidoptera, Zygaenidae] and of *Tricladia* MERCET, 1918 [Hymenoptera, Encyrtidae, Tetracneminae].)

C. & R. FELDER (1874: 9). — KIRBY (1892: 889). — NEAVE (1940: 550). — VIETTE (1951d: 79). — PACLT (1953: 143). — NYE & FLETCHER (1991: 314).

= *Lamelliformia* VIETTE (1951e: 1274): type-species: *Dalaca prytanes* SCHAUS, 1892 by original designation; included species: *tupi* (PFITZNER, 1914), *prytanes* (SCHAUS, 1892); ♂ (1952a: 142). — PACLT (1953: 145); ♂ (1957: 51). — EDWARDS & HOPWOOD (1966: 143). — NYE & FLETCHER (1991: 165); **syn. n.**

Aepytus (Lamelliformia): NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Aepytus (Tricladia): NIELSEN & ROBINSON (1983: 20), syn.: *Aepytus (Pseudophassus)* PFITZNER, 1914, *Aepytus (Parana)* VIETTE, 1950. — ROBINSON & NIELSEN (1984: 17), syn.: *Aepytus (Pseudophassus)* PFITZNER, 1914, *Aepytus (Parana)* VIETTE, 1950.

Cibyra (Lamelliformia): NIELSEN et al. (2000: 843). — GREHAN (2010: 44).

Cibyra (Tricladia): NIELSEN et al. (2000: 843), syn.: *Cibyra (Pseudophassus)* PFITZNER, 1914, *Cibyra (Parana)* VIETTE, 1950. — GREHAN (2010: 49).

Cibyra (Lamelleformia) [sic]: GREHAN (2010: 52).

Remarks. *Lamelliformia* VIETTE, 1951 was described based on the ♂ of *T. prytanes* (SCHAUS, 1892) and *Tricladia* C. & R. FELDER, 1874 on the ♀ of *T. umbrifera* C. & R. FELDER, 1874, which is the same or a closely related species. Species were matched through examination of morphology and mtDNA sequences. All holotype specimens were examined.

Tricladia prytanes (SCHAUS, 1892), **comb. n.**

Dalaca prytanes SCHAUS (1892: 329); Brazil, [Rio de Janeiro], Petrópolis, SCHAUS leg.; coll. SCHAUS; [gen. slide. P. VIETTE no. 91518, type no. 18608]; [USNM]. — BERTKAU (1893: 190). — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1295).

Lamelliformia prytanes: VIETTE (1952a: 143; fig. 7); [syn-] type ♂, GP P. VIETTE no. 2242.

Aepytus (Lamelliformia) prytanes: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Lamelliformia) prytanes: NIELSEN et al. (2000: 843).

Cibyra (Lamelleformia) [sic] *prytanes* [sic]: GREHAN (2010: 52; fig. g appendix).

Remarks. Examination of the holotype places this species in *Tricladia* C. & R. FELDER, 1874, the senior synonym of *Lamelliformia* VIETTE, 1951.

***Tricladia sladeni* (HAMPSON, 1903), comb. n.**

Dalaca sladeni HAMPSON (1903: 260); [holo-]type ♂; Brazil, Mato Grosso, [Santa Anna da Chapada], Chapada, A. ROBERT leg.; [BMNH]. — WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1294).

Aepytus (Hampsoniella) sladeni: VIETTE (1950a: 77; fig. 4 ♂ gen. [error, unidentified species]).

Hampsoniella sladeni: VIETTE (1951c: 95).

Aepytus (Lamelliformia) sladeni: NIELSEN & ROBINSON (1983: 20), syn.: *Aepytus (Lamelliformia) tupi* (PFITZNER, 1914). — ROBINSON & NIELSEN (1984: 17), syn.: *Aepytus (Lamelliformia) tupi* (PFITZNER, 1914).

Cibyra (Lamelliformia) sladeni: NIELSEN et al. (2000: 843), syn.: *Cibyra (Lamelliformia) tupi* (PFITZNER, 1914).

***Tricladia tupi* (PFITZNER, 1914), stat. rev., comb. n.**

Cibyra tupi PFITZNER (1914: 105); Southern Brazil, Sao [RECTE São] Paulo, [Iperó], Ypanema [RECTE Ipanema]; coll. SEITZ; [SMFL].

Hepialus (Cibyra) tupi: PFITZNER (1937: 1293; pl. 99c [♂] dorsal); single specimen in coll. SEITZ.

Lamelliformia tupi: VIETTE (1951e: 1274).

Hepialus tupi: SCHRÖDER (1967: 338); “holotype” [RECTE lectotype] ♂, SMFT 79, GP VIETTE no. 2348. — BIEZANKO (1961a: 8).

Aepytus (Lamelliformia) tupi: NIELSEN & ROBINSON (1983: 20); as synonym of *Aepytus (Lamelliformia) sladeni* (HAMPSON, 1903). — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytus (Lamelliformia) sladeni* (HAMPSON, 1903).

Cibyra (Lamelliformia) tupi: NIELSEN et al. (2000: 843); as synonym of *Cibyra (Lamelliformia) sladeni* (HAMPSON, 1903).

Remarks. The lectotype was examined, and based on genitalic differences it is a species distinct from *T. sladeni* (HAMPSON, 1903).

***Tricladia umbrifera* C. & R. FELDER, 1874, comb. rev.**

C. & R. FELDER (1874: 9; pl. 80, fig. 2 [♀] dorsal); Brasilia [Brazil]. — KIRBY (1892: 889).

Phassus umbrifera: WAGNER & PFITZNER (1911: 19). — PFITZNER (1938: 1300; pl. 185f [♀] dorsal).

Aepytus (Tricladia) umbrifera: NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Tricladia) umbrifera: NIELSEN et al. (2000: 843).

***Vietteogorgopsis* ÖZDIKMEN, 2007**

Type-species: *Paragorgopsis pittionii* VIETTE, 1952 by original designation by VIETTE (1952a: 140) (replacement name).

Paragorgopsis VIETTE (1952a: 140); included species: *pittionii* VIETTE, 1952, *foetterlei* VIETTE, 1952, *schausi* VIETTE, 1952; **praecoc.**: *Paragorgopsis* GIGLIO-TOS, 1893 [Diptera].

PAULT (1957: 52). — EDWARDS & HOPWOOD (1966: 209). — NYE & FLETCHER (1991: 221).

Aepytus (Paragorgopsis): NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopsis): NIELSEN et al. (2000: 842). — GREHAN (2010: 49; fig. e appendix).

Vietteogorgopsis ÖZDIKMEN (2007: 116); replacement name for *Paragorgopsis* VIETTE, 1952. — GREHAN (2010: 47).

***Vietteogorgopsis absyrtus* (SCHAUS, 1892), comb. n.**

Phassus absyrtus SCHAUS (1892: 330); Brazil, [Rio de Janeiro], Petrópolis, SCHAUS leg.; [GP P. VIETTE 91517, type no. 18612],

USNM. — BERTKAU (1893: 190). — WAGNER & PFITZNER (1911: 17). — PFITZNER (1938: 1299). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Remarks. One syntype in the USNM that bears a label “type” is here designated lectotype; it has the following labels: /Petropolis, Brazil/ *Phassus absyrtus* type SCHAUS/ type n°. 18612 USNM/ Coll. Wm. SCHAUS/ Genitalia slide by P. VIETTE USNM 91517/ Photo, det. E. S. NIELSEN 1984/ Photograph on file USNM/. The lectotype was examined, and based on the morphology (especially veins R_{3+4} - R_5 stalked, wing shape) this taxon is placed in *Vietteogorgopsis* ÖZDIKMEN, 2007. The designation is made to permanently stabilise the species’ identification.

***Vietteogorgopsis foetterlei* (VIETTE, 1952)**

Paragorgopsis foetterlei VIETTE (1952a: 141); holotype ♂, Brazil, Rio de Janeiro, Petrópolis, 28. VIII. 1913, J. G. FOETTERLE [leg.]; GP P. VIETTE 1351; NHMW.

Aepytus (Paragorgopsis) foetterlei: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopsis) foetterlei: NIELSEN et al. (2000: 842).

Vietteogorgopsis foetterlei: ÖZDIKMEN (2007: 117).

***Vietteogorgopsis jordani* (VIETTE, 1956)**

Paragorgopsis jordani VIETTE (1956: 377; fig. 3 ♂ gen.); holotype ♂, Brazil, Minas Gerais, Theophilo Ottoni [RECTE Teófilo Ottoni], San Jacintho Valley, 1907-8, F. BIRCH [leg.]; GP P. VIETTE 2329; BMNH.

Aepytus (Paragorgopsis) jordani: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopsis) jordani: NIELSEN et al. (2000: 842).

Vietteogorgopsis jordani: ÖZDIKMEN (2007: 117). — KOÇAK & KOÇAK (2008: 31).

***Vietteogorgopsis katharinae* (PFITZNER, 1914), comb. n.**

Dalaca katharinae PFITZNER (1914: 110); [Brazil, Santa Catarina]; coll. PFITZNER; [SMFL]; ♂ (1937: 1296). — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Dalaca guarani katharinae: SCHRÖDER (1967: 339); “holotype” [RECTE lectotype] [♂], SMFT 98; GP VIETTE no. 2353.

Remarks. The lectotype was examined and, based on morphology (especially veins R_{3+4} - R_5 stalked), this taxon is placed in *Vietteogorgopsis* ÖZDIKMEN, 2007.

***Vietteogorgopsis nigrovenosalis* (VIETTE, 1956)**

Paragorgopsis nigrovenosalis VIETTE (1956: 375; fig. 6 ♂ gen.); holotype ♂, Brazil, Minas Gerais, Aqua [RECTE Água] Suja, x. 1906, E. A. BAER [leg.]; GP P. VIETTE 2990; BMNH.

Aepytus (Paragorgopsis) nigrovenosalis: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopsis) nigrovenosalis: NIELSEN et al. (2000: 842).

Vietteogorgopsis nigrovenosalis: ÖZDIKMEN (2007: 117).

***Vietteogorgopsis pittionii* (VIETTE, 1952)**

Paragorgopsis pittionii VIETTE (1952a: 141; figs. 4 ♂ dorsal, 5 ♂ gen.); holotype ♂, [Brazil], Rio de Janeiro, Petrópolis, 14. XI. 1907, J. G. FOETTERLE [leg.]; GP P. VIETTE 1350; NHMW.

Aepytus (Paragorgopsis) pittionii: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopsis) pittionii: NIELSEN et al. (2000: 842).

Vietteogorgopsis pittionii: ÖZDIKMEN (2007: 117).

***Vietteogorgopsis spitzi* (VIETTE, 1956)**

Paragorgopsis spitzi VIETTE (1956: 375; fig. 2 ♂ gen.); holotype ♂, Brazil, São Paulo, Ypiranga [RECTE Ipiranga], VIII. 1922, R.

SPITZ [leg.]; GP P. VIETTE 2297; BMNH.

Aepytus (Paragorgopis) spitzii: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Paragorgopis) spitzii: NIELSEN et al. (2000: 842).

Vietteogorgopis spitzii: ÖZDIKMEN (2007: 117).

Yleuxas VIETTE, 1951, stat. rev.

Type-species: *Yleuxas bradleyi* VIETTE, 1951, by original designation; monotypic.

VIETTE (1951e: 1280). — PAULT (1953: 145); as synonym of *Aepytus* HERRICH-SCHÄFFER, [1858]. — EDWARDS & HOPWOOD (1966: 316). — NYE & FLETCHER (1991: 323).

Aepytus (Yleuxas): NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Yleuxas): NIELSEN et al. (2000: 843). — GREHAN (2010: 49).

Remarks. *Yleuxas* VIETTE, 1951 was described mainly based on the ♂ genitalia characters, clearly distinct from other genera. Therefore, we re-instate it here as a separate genus.

Yleuxas bradleyi VIETTE, 1951, stat. rev.

Yleuxas bradleyi VIETTE (1951e: 1280; fig. 4 ♂ gen.); holotype ♂, Peru, S. Domingo, Carabaya, 6000 ft., end of wet season, IV. 1901, OCKENDEN [leg.]; GP P. VIETTE no. 2295; BMNH.

Aepytus (Yleuxas) bradleyi: NIELSEN & ROBINSON (1983: 20); as synonym of *Aepytus (Yleuxas) brunnea* (SCHAUS, 1901). — ROBINSON & NIELSEN (1984: 17); as synonym of *Aepytus (Yleuxas) brunnea* (SCHAUS, 1901).

Cibyra (Yleuxas) bradleyi: NIELSEN et al. (2000: 843); as synonym of *Cibyra (Yleuxas) brunnea* SCHAUS, 1901.

Remarks. The ♂ genitalia confirm that this species is not a synonym of *Y. brunnea* (SCHAUS, 1901).

Yleuxas brunnea (SCHAUS, 1901), comb. n.

Cibyra brunnea SCHAUS (1901: 77); Venezuela, Aroa; [GP VIETTE no. 91525, type no. 18609]; [USNM].

Hepialus (Cibyra) brunnea: PFITZNER (1937: 1293).

Aepytus (Yleuxas) brunnea: NIELSEN & ROBINSON (1983: 20), syn.: *Aepytus (Yleuxas) bradleyi* (VIETTE, 1951). — ROBINSON & NIELSEN (1984: 17), syn.: *Aepytus (Yleuxas) bradleyi* (VIETTE, 1951).

Cibyra (Yleuxas) brunnea: NIELSEN et al. (2000: 843), syn.: *Cibyra (Yleuxas) bradleyi* (VIETTE, 1951).

Remarks. One syntype ♂ that bears a label “type” is here designated lectotype; it has the following labels: /Aroa, Venezuela. / *Cibyra brunnea* type SCHAUS/ type n°. 18609 USNM/ Coll. Wm. SCHAUS/ Genitalia slide by P. VIETTE USNM 91525/ Photo, det. E. S. NIELSEN 1984. The lectotype was examined and it is different from *Y. bradleyi* VIETTE, 1951, also examined. The designation is made to permanently stabilise the species’ identification.

Taxa incertae sedis

Cibyra (Aepytus) petropoliensis (VIETTE, 1952)

Aepytus petropoliensis VIETTE (1952a: 140; fig. 1 ♀ dorsal); holotype ♀, Brazil, Petrópolis, 25. III. 1903, J. G. FOETTERLE [leg.]; specimen without abdomen; NHMW.

Aepytus (Aepytus) petropoliensis: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Aepytus) petropoliensis: NIELSEN et al. (2000: 842).

Remarks. The holotype was examined, and the wings markings were found to be distinct from all known genera. A genitalia dissection was not possible because the abdomen is lacking.

Cibyra (Gymelloxes) paropus (DRUCE, 1890)

Hepialus paropus DRUCE (1890: 508); Ecuador, Sarayacu, BUCKLEY [leg.]; coll. DRUCE; [GP P. VIETTE no. 2026]; [BMNH]. — BERTKAU (1891: 196). — KIRBY (1892: 884). — WAGNER & PFITZNER (1911: 9). — ZUKOWSKI (1954: 93).

Hepialus (Hepialus) paropus: PFITZNER (1937: 1291).

Aepytus (Gymelloxes) paropus: NIELSEN & ROBINSON (1983: 19). — ROBINSON & NIELSEN (1984: 17).

Cibyra (Gymelloxes) paropus: NIELSEN et al. (2000: 842).

Remarks. The holotype was examined and found to show no obvious similarity to the type species of *Gymelloxes* VIETTE, 1952 and to any other of the described genera.

Dalaca cocama PFITZNER, 1914

Dalaca (Triodia) cocama PFITZNER (1914: 110); Peru, Huanca-bamba, 6400'; coll. PFITZNER; [SMFL].

Dalaca cocama: PFITZNER (1937: 1296). — VIETTE (1951e: 1282). — ZUKOWSKI (1954: 93). — SCHRÖDER (1967: 339); “holotype” [RECTE lectotype] [♀], SMFT 95. — NIELSEN & ROBINSON (1983: 20), syn.: *Dalaca nannophyes* PFITZNER, 1914. — ROBINSON & NIELSEN (1984: 17), syn.: *Dalaca nannophyes* PFITZNER, 1914. — NIELSEN et al. (2000: 843), syn.: *Dalaca nannophyes* PFITZNER, 1914.

= *Dalaca (Triodia) nannophyes* PFITZNER (1914: 110); [Colombia], Sosomoco, 800 m, FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19).

Dalaca (Triodia) nannophyes: PFITZNER (1938: 1297); single specimen [= holotype?], [Colombia], Sosomoco, 800 m, FASSL leg; praecoc.: PFITZNER, 1914 [Hepialidae].

Dalaca cocama nannophyes: SCHRÖDER (1967: 339); “holotype” [RECTE lectotype] ♂, SMFT 96, GP VIETTE no. 2354.

Dalaca nannophyes: NIELSEN & ROBINSON (1983: 20); as synonym of *Dalaca cocama* PFITZNER, 1914. — ROBINSON & NIELSEN (1984: 17); as synonym of *Dalaca cocama* PFITZNER, 1914. — NIELSEN et al. (2000: 843); as synonym of *Dalaca cocama* PFITZNER, 1914.

Remarks. Placing the taxa *cocama* PFITZNER, 1914, *nannophyes* PFITZNER, 1914 and *nannophyes* PFITZNER, 1938 [identical with *nannophyes* PFITZNER, 1914?] into a genus requires further studies. Therefore, for the time being, we prefer to retain the last (provisional) systematic position as defined by NIELSEN & ROBINSON (1983).

Dalaca cuprifera PFITZNER, 1914

Dalaca cuprifera PFITZNER (1914: 105); Peru; ex coll. STAUDINGER; [SMFL]; ♂ (1937: 1294; pl. 99f [♀] dorsal). — ZUKOWSKI (1954: 93). — SCHRÖDER (1967: 339); “holotype” [RECTE lectotype] [♀], SMFT 88. — NIELSEN & ROBINSON (1983: 20). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. It is difficult to match the lectotype (♀) to any known ♂ and, anyway, placing it into any described genus depends on further investigation. So, we keep the systematic position chosen by NIELSEN & ROBINSON (1983).

Dalaca manoa PFITZNER, 1914

Dalaca manoa PFITZNER (1914: 105); single ♀ [= holotype], Ostkolumbien (East Colombia), Villavicencio, 450 m, II. 1911, FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19). — PFITZNER (1937: 1294; pl. 99d [♀] dorsal). — SCHRÖDER (1967: 340); holotype ♀, SMFT 89. — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. It is difficult to match the holotype (♀) to any known ♂ and, anyway, placing it into any described genus depends on further investigation (possibly *Pfitzneriana*?). So, we keep the systematic position chosen by NIELSEN & ROBINSON (1983).

***Dalaca mummia* SCHAUS, 1892**

Dalaca mummia SCHAUS (1892: 330); holotype ♀, Brazil, [Rio de Janeiro], Petrópolis, SCHAUS leg.; coll. SCHAUS; [GP P. VIETTE no. 91520, type no. 18606]; USNM. — BERTKAU (1893: 190). — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Dalaca mummea [sic] WAGNER & PFITZNER (1911: 14). — PFITZNER (1937: 1295).

Remarks. It is difficult to match the holotype (♀) to any known ♂ and, anyway, placing it into any described genus depends on further investigation. So, we keep the systematic position chosen by NIELSEN & ROBINSON (1983).

***Dalaca niepelti* PFITZNER, 1914**

Dalaca niepelti PFITZNER (1914: 59; pl. 11, figs. 14 ♂ dorsal, 15 ♀ dorsal); Ecuador, Macas; coll. NIEPELT; [BMNH]. — STRAND (1927: 42). — PFITZNER (1937: 1296). — ZUKOWSKI (1954: 94). — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. It is difficult to match the holotype (♀) to any known ♂ and, anyway, placing it into any described genus depends on further investigation. So, we keep the systematic position chosen by NIELSEN & ROBINSON (1983).

***Dalaca usaque* PFITZNER, 1914**

Dalaca usaque PFITZNER (1914: 105); Colombia, Muzo, 700 m, FASSL [leg.]; coll. PFITZNER; [SMFL]; ♂ (1937: 1294; pl. 99f [♂] dorsal). — SCHRÖDER (1967: 340); “holotype” [recte lectotype: PFITZNER described the taxon based on an uncertain number of specimens, so a type designation creates a lectotype] ♂, SMFT 82; East Gramal near Muzo; GP VIETTE no. 2351. — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. The holotype ♂ does not fit into any known genus, so we prefer to retain the last (provisional) systematic position as defined by NIELSEN & ROBINSON (1983).

***Dalaca vibicata* PFITZNER, 1914**

Dalaca vibicata PFITZNER (1914: 105); [Colombia], Sosomoco, FASSL [leg.]; coll. PFITZNER; [SMFL]. — FASSL (1918: 19). — PFITZNER (1937: 1294; pl. 99c [♂] dorsal). — SCHRÖDER (1967: 340); “holotype” [recte lectotype, see *usaque* PFITZNER, 1914, similar case] ♂, SMFT 86; III. 1911; GP VIETTE no. 2350. — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. The holotype ♂ does not fit into any known genus, so we prefer to retain the last (provisional) systematic position as defined by NIELSEN & ROBINSON (1983).

“*Hepialus* sp.”

Hepialus sp.: KOEBELE (1924: 56–59, 68). — SWEZEY (1925: 376). — GARA & ONORE (1989: 142; fig. 115 larva).

Remarks. This reference of an unidentified species in definitively an incorrect genus exists, so until we know which species the authors were dealing with, we keep this data here. It cannot, of course, be included in the checklist.

***Phassus costaricensis* DRUCE, 1887**

Phassus costaricensis DRUCE (1887: 234; pl. 24, fig. 4 [♀] dorsal); single specimen [= holotype], Costa Rica, VAN PATTEN [leg.]; BMNH. — KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 17). — PFITZNER (1938: 1300; pl. 185c [♀] dorsal). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Dalaca costaricensis: FORBES (1942: 406); as synonym of *Dalaca assa* DRUCE, 1887.

Remarks. The holotype was examined, and it was found to lack a wing pattern similar to that of the type-species of the genus *Phassus* WALKER, 1856. The holotypes of *costaricensis* and *assa* appear to be different, so FORBES’ (1942) synonymisation requires further studies. It is difficult to match the holotype of *costaricensis* (♀) to any known ♂ and, anyway, placing it into any described genus depends on further investigation. So, we keep the generic association chosen by NIELSEN et al. (2000); the species was moved to the *incertae sedis* position by us.

***Phassus eldorado* PFITZNER, 1906**

PFITZNER (1906: 276); single ♂ [= holotype], Venezuela, Mérida; coll. PFITZNER; [SMFL]. — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1298; pl. 99g [♂] dorsal). — SCHRÖDER (1967: 342); holotype ♂, SMFT 48; GP VIETTE no. 2355. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Remarks. The holotype was examined, and it was found to lack a wing pattern similar to that of the type-species of the genus *Phassus* WALKER, 1856. Placing the taxon into any genus depends on further investigation, so we keep the generic association chosen by NIELSEN et al. (2000); the species was moved to the *incertae sedis* position by us.

***Phassus guianensis* SCHAUS, 1940**

SCHAUS (1940: 83, 88); [holo-]type [♂], British Guiana, Karta; type no. 34749; USNM. — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Remarks. The holotype was examined, and it was found to lack a wing pattern similar to that of the type-species of the genus *Phassus* WALKER, 1856. Placing the taxon into any genus depends on further investigation, so we keep the generic association chosen by NIELSEN et al. (2000); the species was moved to the *incertae sedis* position by us.

***Phassus pretiosus* (HERRICH-SCHÄFFER, [1856])**

Epialus [sic] *pretiosus* HERRICH-SCHÄFFER ([1856]: cover; pl. 88, fig. 505); Brazil; ♂ ([1856]: 5; fig. 505 [♂] dorsal); ♂ ([1858]: 57, 84; fig. 505), syn.: *plusia* BOISDUVAL, *in litt.*

= *Epialus plusia* HERRICH-SCHÄFFER ([1856]: 57); as synonym of *Epialus* [sic] *pretiosus* HERRICH-SCHÄFFER, [1856]

Hepialus pretiosus: KIRBY (1892: 884). — WAGNER & PFITZNER (1911: 9).

Hepialus (Hepialus) pretiosus: PFITZNER (1937: 1291).

Phassus pretiosus: NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Remarks. The holotype was not found, but according to HERRICH-SCHÄFFER’s figure, it appears to lack a wing pattern similar to that of the type-species of the genus *Phassus* WALKER, 1856. Placing the taxon into any genus depends on further investigation, so we keep the generic association chosen by NIELSEN et al. (2000); the species was moved to the *incertae sedis* position by us.

***Phassus smithi* DRUCE, 1889**

DRUCE (1889: 92); single specimen [= holotype], Mexico, Atoyac, Vera Cruz, v. 1888, H. H. SMITH [leg.]; [BMNH]. — KIRBY (1892: 890). — DRUCE (1898: 452; pl. 89, fig. 5 [♀] dorsal). — WAGNER & PFITZNER (1911: 18). — PFITZNER (1938: 1300; pl. 185d [♀] dorsal). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 16). — NIELSEN et al. (2000: 841).

Remarks. The holotype was examined, and it was found to lack a wing pattern similar to that of the type-species of the genus *Phassus* WALKER, 1856. It is difficult to match the holotype (♀) to any known ♂ and placing it into any described genus depends on further investigation. So, we keep the generic association chosen by

NIELSEN et al. (2000); the species was moved to the *incertae sedis* position by us.

***Phassus transversus* WALKER, 1856**

WALKER (1856: 1567); [Brazil], Rio de Janeiro; coll. FRY [not found in Oxford University Museum, UK]. — KIRBY (1892: 890). — WAGNER & PFITZNER (1911: 19). — NIELSEN & ROBINSON (1983: 18). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 841).

Remarks. The holotype was not examined, but according to geographical distribution and wing description, it was found to evidently lack any close relationship to the type-species of the genus *Phassus* WALKER, 1856. Placing the taxon into any genus depends on further investigation, so we keep the generic association chosen by NIELSEN et al. (2000); the species was moved to the *incertae sedis* position by us. Possibly, the taxon belongs to the genus *Tricladia*.

Sedis novum

***Acrolophus tapuja* (PFITZNER, 1914) (Tineidae, new family and genus combination)**

Dalaca tapuja PFITZNER (1914: 110); Southern Brazil, Leopoldina; coll. SEITZ; [SMFL]. — PFITZNER (1937: 1296; pl. 99e dorsal). — NIELSEN & ROBINSON (1983: 21). — ROBINSON & NIELSEN (1984: 17). — NIELSEN et al. (2000: 844).

Remarks. The holotype was examined and it is a tineid, to be placed within the genus *Acrolophus* POEY, 1832 (V. O. BECKER pers. comm.).

Checklist

(Arrangement alphabetically as in catalogue; without synonyms.)

***Aepytyus* HERRICH-SCHÄFFER, [1856] stat. rev.**

Aepytyus biedermanni (VIETTE, 1950), **comb. rev.**
Aepytyus exclamans (HERRICH-SCHÄFFER, [1854]), **comb. rev.**
Aepytyus guarani (PFITZNER, 1914), **comb. n.**

***Alloaepytyus* VIETTE, 1951, stat. rev.**

Alloaepytyus tesselloides (SCHAUS, 1901), **comb. rev.**

***Andeabatis* NIELSEN & ROBINSON, 1983**

Andeabatis chilensis (URETA, 1951)

***Aplatissa* VIETTE, 1953**

Aplatissa michaelis (PFITZNER, 1914)
Aplatissa strangoides VIETTE, 1953

***Blanchardinella* NIELSEN, ROBINSON & WAGNER, 2000**

Blanchardinella venosus (BLANCHARD, 1852)

***Calada* NIELSEN & ROBINSON, 1983**

Calada fuegensis NIELSEN & ROBINSON, 1983
Calada migueli NIELSEN & ROBINSON, 1983

***Callipielus* BUTLER, 1882**

Callipielus arenosus BUTLER, 1882
Callipielus argentata URETA, 1957
Callipielus digitata ROBINSON, 1977
Callipielus fumosa NIELSEN & ROBINSON, 1983
Callipielus gentilii NIELSEN & ROBINSON, 1983
Callipielus izquierdoi (URETA, 1957)
Callipielus krahmeri NIELSEN & ROBINSON, 1983
Callipielus perforata NIELSEN & ROBINSON, 1983
Callipielus salasi ROBINSON, 1977
Callipielus vulgaris NIELSEN & ROBINSON, 1983

***Cibyra* WALKER, 1856**

Cibyra danieli (VIETTE, 1961)
Cibyra dorita SCHAUS, 1901
Cibyra ferruginosa WALKER, 1856
Cibyra forsteri (VIETTE, 1961)
Cibyra monoargenteus (VIETTE, 1951)
Cibyra munona (SCHAUS, 1929)
Cibyra oreas (SCHAUS, 1892), **comb. rev.**
Cibyra pluriargenteus (VIETTE, 1956)
Cibyra schausi (VIETTE, 1952), **comb. rev.**
Cibyra stigmatica (PFITZNER, 1937), **comb. n.**
Cibyra verresi (SCHAUS, 1929)
Cibyra yungas (VIETTE, 1961)
Cibyra zischkai (VIETTE, 1961)

***Dalaca* WALKER, 1856**

Dalaca chiliensis (VIETTE, 1950)
Dalaca crocatus (URETA, 1956)
Dalaca laminata NIELSEN & ROBINSON, 1983
Dalaca nigricornis WALKER, 1856
Dalaca pallens (BLANCHARD, 1852)
Dalaca parafuscus NIELSEN, ROBINSON & WAGNER, 2000
Dalaca patriciae NIELSEN & ROBINSON, 1983
Dalaca postvariabilis NIELSEN & ROBINSON, 1983
Dalaca quadricornis NIELSEN & ROBINSON, 1983
Dalaca variabilis (VIETTE, 1950)

***Druceiella* VIETTE, 1949**

Druceiella amazonensis VIETTE, 1950
Druceiella basirubra (SCHAUS, 1901)
Druceiella metellus (DRUCE, 1890)
Druceiella momus (DRUCE, 1890)

***Gymelloxes* VIETTE, 1952, stat. rev.**

Gymelloxes prosopus (DRUCE, 1901), **comb. n.**
Gymelloxes terea (SCHAUS, 1892), **comb. rev.**
Gymelloxes trilinearis (PFITZNER, 1914), **comb. rev.**

***Hampsoniella* VIETTE, 1950, stat. rev.**

Hampsoniella assa (DRUCE, 1887), **comb. rev.**
Hampsoniella equatorialis (VIETTE, 1950), **comb. rev.**

***Hepialyxodes* VIETTE, 1951, stat. rev.**

Hepialyxodes rileyi VIETTE, 1951, **comb. rev.**

***Parapielus* VIETTE, 1949**

Parapielus heimlichii (URETA, 1956)
Parapielus luteicornis (BERG, 1882)
Parapielus oberthuri (VIETTE, 1951)
Parapielus reedi (URETA, 1957)

***Pfitzneriana* VIETTE, 1952**

Pfitzneriana allura VIETTE, 1961
Pfitzneriana obliquestrigata (STRAND, 1912), **comb. n.**
Pfitzneriana olivescens (PFITZNER, 1914)
Pfitzneriana vogli VIETTE, 1952

***Pfitzneriella* VIETTE, 1951**

Pfitzneriella lucicola (MAASSEN, 1890)
Pfitzneriella monticola (MAASSEN, 1890)
Pfitzneriella remota (PFITZNER, 1906)
Pfitzneriella similis (ZUKOWSKI, 1954)

Phassus WALKER, 1856

- Phassus aurigenus* PFITZNER, 1914
Phassus basirei SCHAUS, 1890
Phassus championi DRUCE, 1887
Phassus chrysodidyma DYAR, 1915
Phassus exclamationis PFITZNER, 1938
Phassus huebneri (GEYER, [1838])
Phassus marcius DRUCE, 1892
Phassus n-signatus WEYMER, 1907
Phassus phalerus DRUCE, 1887
Phassus pharus (DRUCE, 1887)
Phassus rosulentus WEYMER, 1907
Phassus triangularis EDWARDS, 1885

Phialuse VIETTE, 1961

- Phialuse palmar* VIETTE, 1961

Philoenia KIRBY, 1892, **stat. rev.**

- Philoenia brasiliensis* VIETTE, 1952, **comb. rev.**
Philoenia fasslii (PFITZNER, 1914), **comb. rev.**
Philoenia guyanensis (VIETTE, 1951), **comb. rev.**
Philoenia indicata (STRAND, 1912), **comb. n.**
Philoenia lagopus (MÖSCHLER, 1877), **comb. rev.**
Philoenia saguanmachica (PFITZNER, 1914), **comb. rev.**
Philoenia thisbe (DRUCE, 1901), **comb. n.**

Pseudodalaca VIETTE, 1950, **stat. rev.**

- Pseudodalaca gugelmanni* (VIETTE, 1950), **comb. rev.**
Pseudodalaca mexicanensis VIETTE, 1953, **comb. rev.**
Pseudodalaca sarta (SCHAUS, 1894), **comb. n.**

Pseudophassus PFITZNER, 1914, **stat. rev.**

- Pseudophassus mahagoniatus* PFITZNER, 1914, **comb. rev.**
Pseudophassus philipponi (VIETTE, 1950), **comb. n.**

Pseudophilaenia VIETTE, 1951, **stat. rev.**

- Pseudophilaenia omagua* (PFITZNER, 1937), **comb. rev.**

Puermytrans VIETTE, 1951

- Puermytrans chiliensis* VIETTE, 1951

Roseala VIETTE, 1950

- Roseala tessellatus* (HERRICH-SCHÄFFER, [1854]), **comb. n.**

Schaefferiana VIETTE, 1950, **stat. rev.**

- Schaefferiana epigramma* (HERRICH-SCHÄFFER, [1854]), **comb. rev.**
Schaefferiana simplex VIETTE, 1956, **comb. rev.**

Schausiana VIETTE, 1950

- Schausiana trojesa* (SCHAUS, 1901)

Trichophassus LE CERF, 1919

- Trichophassus giganteus* (HERRICH-SCHÄFFER, [1853])

Tricladia C. & R. FELDER, 1874, **stat. rev.**

- Tricladia prytanes* (SCHAUS, 1892), **comb. n.**
Tricladia sladeni (HAMPSON, 1903), **comb. n.**
Tricladia tupi (PFITZNER, 1914), **stat. rev., comb. n.**
Tricladia umbrifera C. & R. FELDER, 1874, **comb. rev.**

Vietteogorgopis ÖZDIKMEŇ, 2007

- Vietteogorgopis absyrtus* (SCHAUS, 1892), **comb. n.**
Vietteogorgopis foetterlei (VIETTE, 1952)
Vietteogorgopis jordani (VIETTE, 1956)

Vietteogorgopis katharinae (PFITZNER, 1914), **comb. n.**

Vietteogorgopis nigrovenosalis (VIETTE, 1956)

Vietteogorgopis pittionii (VIETTE, 1952)

Vietteogorgopis spitzi (VIETTE, 1956)

Yleuxas VIETTE, 1951, **stat. rev.**

Yleuxas bradleyi VIETTE, 1951, **stat. rev.**

Yleuxas brunnea (SCHAUS, 1901), **comb. n.**

Taxa incertae sedis

Cibyra (Aepytus) petropolisensis (VIETTE, 1952)

Cibyra (Gymelloxes) paropus (DRUCE, 1890)

Dalaca cocama PFITZNER, 1914

Dalaca cuprifera PFITZNER, 1914

Dalaca manoa PFITZNER, 1914

Dalaca mummia SCHAUS, 1892

Dalaca niepelti PFITZNER, 1914

Dalaca usaque PFITZNER, 1914

Dalaca vibicata PFITZNER, 1914

Phassus costaricensis DRUCE, 1887

Phassus eldorado PFITZNER, 1906

Phassus guianensis SCHAUS, 1940

Phassus pretiosus (HERRICH-SCHÄFFER, [1856])

Phassus smithi DRUCE, 1889

Phassus transversus WALKER, 1856

Sedis novum: family transfer to Tineidae

Acrolophus tapuja (PFITZNER, 1914), **comb. n.**

References

- AGUIAR-MENEZES, E. L., MENEZES, E. B., CASSINO, P. C. R., & SOARES, M. A. (2002): Passion fruit. — Pp. 360–390 *in*: PENA, J., SHARP, J., & WYSOKI, M. (eds.), Tropical fruit pests and pollinators: biology, economic importance, natural enemies and control. — Trowbridge (CABI), 430 pp.
- AGUILERA P., A., ELLENA D., M., SEGUEL B., I., MONTENEGRO B., A., MARTÍN A., J. S., TORRALBO B., L., & REBOLLEDO R., R. (2009): Nuevos registros de insectos fitófagos asociados a murta o murtila (*Ugni molinae* TURCZ.) em Chile. — IDESIA, Arica (Chile), 27 (2): 33–37.
- ANDRADE, E. N. (1928): Contribuição para o estudo da entomologia florestal paulista. — Boletim de Agricultura, São Paulo, 29 (7/8): 446–453.
- ARGUEDAS, M. (2004): Problemas fitosanitarios de la melina (*Gmelina arborea* (ROXB.)) em Costa Rica. Nota Técnica. — Kurú: Revista Forestal, Cartago (Costa Rica), 1 (2): 1–9.
- (2006): Clasificación de tipos de daños producidos por insectos forestales. Segunda parte. — Kurú: Revista Forestal, Cartago (Costa Rica), 3 (9): 1–8.
- (2007): Plagas y enfermedades forestales em Costa Rica. — Kurú: Revista Forestal, Cartago (Costa Rica), 4 (11/12): vi + 69 pp.
- , & ESPINOZA, D. (2007): Problemas fitosanitarios del jaúl (*Alnus acuminata* KUNTH) em Costa Rica. — Kurú: Revista Forestal, Cartago (Costa Rica), 4 (10): 1–10.
- AURIVILLIUS, P. O. C. (1884): Lepidoptera. — *In*: MAYER, P., & GIESBRECHT, W. (ed.), Zoologischer Jahresbericht für 1883. Herausgegeben von der Zoologischen Station zu Neapel. II. Abtheilung: Arthropoda. — Leipzig (W. Engelmann), [iv] + 585 pp.
- BARRETT, O. W. (1900): Some notes on “The Cambridge Natural history, vol. VI”. — Canadian Entomologist, Ottawa, 32 (8): 234–236.

- BERG, C. (1882): Farrago Lepidopterologia. Contribuciones al estudio de la fauna Argentina y países limítrofes. — Anales de la Sociedad Científica Argentina, Buenos Aires, **13**: 164-184, 213-223, 257-279.
- BERTKAU, P. (1891): Bericht über die wissenschaftlichen Leistungen im Gebiete der Entomologie während des Jahres 1890. — Archiv für Naturgeschichte, Berlin, **57** (2): 1-343.
- (1893): Bericht über die wissenschaftlichen Leistungen im Gebiete der Entomologie während des Jahres 1892. — Archiv für Naturgeschichte, Berlin, **59** (2): 1-348.
- BIEZANKO, C. M. (1961a): Castniidae, Zygaenidae, Dalceridae, Eucleidae, Megalopygidae, Cossidae et Hepialidae da Zona Sueste do Rio Grande do Sul (Contribuição ao conhecimento da fisiografia do Rio Grande do Sul). — Arquivos de Entomologia, Pelotas, (A) **14**: 1-14.
- (1961b): Castniidae, Zygaenidae, Dalceridae, Eucleidae, Megalopygidae, Cossidae et Hepialidae da Zona Missioneira do Rio Grande do Sul (Contribuição ao conhecimento da fisiografia do Rio Grande do Sul). — Arquivos de Entomologia, Pelotas, (B) **14**: 1-8.
- , RUFFINELLI, A., & CARBONELL, C. S. (1957): Lepidoptera del Uruguay. Lista anotada de especies. — Revista de la Facultad de Agronomía, Montevideo, **46**: 1-152.
- BLANCHARD, E. (1852-1854): Fauna Chilena. Insectos. Orden VI Lepidopteros. — In: GAY, C. (ed.), Historia física y política de Chile según documentos adquiridos en esta república durante doce años de residencia en ella y publicada bajo los auspicios del Supremo Gobierno. Zoología. Vol. 7. — Paris (C. Gay), pp. 1-112 (1852), 7 pls. (1854).
- BOURGOGNE, J. (1949): Un type nouveau d'appareil génital femelle chez les Lépidoptères. — Annales de la Société entomologique de France, Paris, **115**: 69-80.
- BRYK, F. (1944): Über die Schmetterlingsausbeute der schwedischen wissenschaftlichen Expedition nach Patagonien 1932-1934. Report Number 11 of the Ljungner Expedition 1932-1934. — Arkiv för Zoologi, Stockholm, **36A** (3): 1-30, 2 pls.
- BRIQUELOT, A. (1956): Sobre a ocorrência de um macrolepidóptero em eucaliptus. — Companhia Siderúrgica Belgo Mineira, Departamento de Terras, Matas e Carvão, Serviço Florestal, Boletim Técnico, Monlevade (MG, Brazil), 37 pp.
- BUTLER, A. G. (1882): Heterocerous Lepidoptera collected in Chili by Thomas EDMONDS, Esq. Part I. Sphingines and Bombyces. — Transactions of the Entomological Society of London, London, **1882**: 1-30.
- BUZZI, Z. J. (1994): Coletânea de nomes populares de insetos do Brasil. — Universidade Federal do Paraná, Departamento de Zoologia, Curitiba, [v] + 230 pp.
- (2009): Nomes populares de insetos e ácaros do Brasil. — Curitiba (Ed. UFPR), 629 pp.
- CABALLERO V., C. (1955): La cuncunilla en los pastos del Sur. — Agricultura y Ganadería, Santiago (Chile), **1** (3): 7-8.
- CAMOUSSEIGHT M., A. (1980): Catálogo de los tipos de insecta depositados en la colección del Museo Nacional de Historia Natural (Santiago, Chile). — Publicación ocasional. Museo Nacional de Historia Natural, Santiago (Chile), **32**: 3-46, 1 map.
- CARRILLO, R. (1974): Eficacia de diversos insecticidas en el combate de cuncunillas negras en empastadas. — Agro Sur, Valdivia (Chile), **1** (2): 45-50.
- CENGICANA (CENTRO GUATEMALTECO DE INVESTIGACIÓN Y CAPACITACIÓN DE LA CAÑA DE AZÚCAR) (ed.) (2007): Eventos históricos y logros 1992-2007. — Guatemala City (Guatemala), 85 pp.
- (2008): Informe anual 2006-2007, Guatemala City (Guatemala), 100 pp.
- CISTERNAS A., E. (1989): Caracterización morfológica de insectos asociados al suelo en el Sur de Chile. — Instituto de Investigaciones Agropecuarias, Estación experimental Remehue, Boletín técnico, Osorno (Chile), **150**: 1-19.
- (1992): Biología y control de insectos plagas em praderas. — In: Seminario manejo de praderas permanentes, Serie Remehue, Osorno (Chile), **31**: 87-117.
- (2000): La cuncunilla negra. — Informativo Remehue, Osorno (Chile), **16**: 1-4.
- , FRANCE I., A., & GERDING P., M. (2007): Control de cuncunilla negra entomopatígeno. — INIA Tierra adentro, Santiago (Chile), **9-10**: 42-43.
- , GERDING G., M., & FRANCE L., A. (2003): Uso del hongo *Beauveria bassiana*: ¿Cómo controlar la cuncunilla negra de las praderas sin usar insecticidas? — Informativo Agropecuario Bioleche INIA-Quilamapu, Quilamapu (Chile), **16** (1): 51-52.
- DAPOTO, G., GIGANTI, H., GENTILI, M., & BONDONI, M. (2003): Lepidópteros de los bosques nativos del Departamento Aluminé (Neuquén-Argentina). II Contribución. — Bosque, Valdivia (Chile), **24** (1): 95-112.
- DAY, M. D., WILEY, C. J., PLAYFORD, J., & ZALUCKI, M. P. (2003): *Lantana*: current management status and future prospects. — ACIAR (Australian Centre for International Agricultural Research), Canberra, Monograph **102**, vii + 128 pp.
- DEVOTTO, L., CISTERNAS, E., CARRILLO, R., & GERDING, M. (2008): Non-target effects of *Dalaca pallens* BLANCHARD control examined through principal response curves: a guild approach in Southern Chile. — Chilean Journal of Agricultural Research, Chillán (Chile), **68** (3): 228-237.
- , ——, GERDING, M., & CARRILLO, R. (2007): Response of grassland soil arthropod community to biological and conventional control of a native moth: using *Beauveria bassiana* and lambda-cyhalothrin for *Dalaca pallens* (Lepidoptera: Hepialidae) suppression. — BioControl, Dordrecht (NL), **52**: 507-531.
- DRUCE, H. (1881-1900): Heterocera. — In: GODMAN, F. C., & SALVIN, O. (eds.), Biología Centrali-Americana, Zoology. Insecta, Lepidoptera-Heterocera, vols. I: xxxi + 490 pp., II: 622 pp., III: 101 col. pls. — London.
- (1889): Descriptions of new species of Lepidoptera, chiefly from Central America. — Annals and Magazine of Natural History, London, (6) **4**: 77-94.
- (1890): Descriptions of new species of Lepidoptera Heterocera from Central and South America. — Proceedings of the Zoological Society of London, London, **1890**: 493-520.
- (1892): Description of a new genus and some species of Heterocera from Central America. — Annals and Magazine of Natural History, London, (6) **9**: 275-279.
- (1901): Descriptions of some new species of Lepidoptera from East Africa and tropical America. — Annals and Magazine of Natural History, London, (7) **7**: 432-444.
- DUGDALE, J. S. (1994): Hepialidae (Insecta: Lepidoptera). — In: DUVAL, C. T. (ed.), Fauna of New Zealand/Ko te Aitanga Pepeke o Aotearoa, vol. **30**. — Lincoln, Canterbury (New Zealand) (Manaaki Whenua), 164 pp., 349 + 25 figs., 27 + 5 maps.
- DUMBLETON, L. J. (1966): Genitalia, classification and zoogeography of the New Zealand Hepialidae (Lepidoptera). — New Zealand Journal of Science, Wellington, **9** (4): 920-981.
- DURÁN M., L. (1976): Problemas de la entomología agrícola em Chile Austral. — Agro Sur, Valdivia (Chile), **4** (2): 119-127.
- DYAR, H. G. (1915a): New American Lepidoptera chiefly from Mexico. — Insector Inscitiae Menstruus, Washington D.C., **3**: 79-85.

- (1915b): Report on the Lepidoptera of the Smithsonian Biological Survey of the Panama Zone. — Proceedings of the United States National Museum, Washington D.C., **47**: 139–350.
- (1917): Description of some lepidopterous larvae from Mexico. — *Insector Inscitiae Menstruus*, Washington D.C., **5** (1–3): 128–132.
- EDWARDS, H. (1885): Notes on Mexican Lepidoptera, with descriptions of new species. — *Entomologica Americana*, New York, **1** (7): 128–129.
- EDWARDS, M. A., & HOPWOOD, A. T. (1966): *Nomenclator Zoologicus*, vol. 6. — Zoological Society of London, London, x + 329 pp.
- , & MANLY, P., & TOBIAS, M. (1996): *Nomenclator Zoologicus*, vol. 9. — Zoological Society of London, London, iv + 747 pp.
- , & VEVERS, H. G. (1975): *Nomenclator Zoologicus*, vol. 7. — Zoological Society of London, London, iii + 374 pp.
- ENDERLEIN, G. (1912): Die Insekten des Antarkto-Archiplatea-Gebietes (Feuerland, Falkland-Inseln, Süd-Georgien). 20. Beitrag zur Kenntnis der Antarktischen Fauna. — *Kungliga Svenska Vetenskapsakademiens Handlingar*, Stockholm, **8** (3): 170 pp. + 4 pls.
- FASSL, A. H. (1918): Verzeichnis neu beschriebener Schmetterlingsformen und Jugendzustände tropischer Lepidopteren von meiner Columbien-Reise (1908–1912). — *Entomologische Rundschau*, Stuttgart, **33**: 1–31.
- FELDER, C., & FELDER, R. (1874): Heft 4 (partim), Heterocera, Bombyces & Sphinges, plates 75–107. — *In*: FELDER, C., FELDER, R., & ROGENHOFER, A. F. (eds.) ([1864] 1865–1875), *Reise der österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von WÜLLERSTORF-URBAIR*. Zoologischer Theil, Zweiter Band, Zweite Abtheilung: Lepidoptera, Atlas [= butterflies plates + Hefte 4 + 5 with all legends and explanations]. — Vienna (K.-k. Hof- und Staatsdruckerei), pp. 1–9, 1–10, 1–20, pls. 1–140.
- FOETTERLE, J. G. (1903): Descrição de Lepidopteros novos do Brazil. — *Revista do Museu Paulista*, São Paulo, **5**: 618–653, 1 pl.
- FORBES, W. T. M. (1942): The Lepidoptera of Barro Colorado Island, Panama, no. 2. — *Bulletin of the Museum of Comparative Zoology at Harvard College*, Cambridge (Mass.), **90** (2): 261–406, pls. 9–16.
- GAEDE, M. (1930): 24. Familie: Hepialidae. — Pp. 553–559, pl. 80 *in*: SEITZ, A. (ed.) (1926–1930), *Die Gross-Schmetterlinge der Erde*, **14**. Band. Die Afrikanischen Spinner und Schwärmer. — Stuttgart (A. Kernen), vii + 597 + vii pp., pls. 1–80.
- GARA, R. I., & ONORE, G. (1989): *Entomologia forestal*. — Proyecto DINAFAID, Quito (Ecuador), 267 pp.
- GERSTAECKER, A. (1857): Bericht die wissenschaftlichen Leistungen im Gebiete der Entomologie während des Jahres 1856. — *Archiv für Naturgeschichte*, Berlin, **23** (2): 273–486.
- GEYER, C. ([1838]): pl. [53] with legend, *in*: HÜBNER, J. ([1827–1838] [dated according to HEMMING 1937]), *Sammlung exotischer Schmettlinge* [sic], vol. 3. — Augsburg (J. Hübner), 53 pls.
- GIGANTI, H., DAPOTO, G., & GENTILI, P. (1994): Lepidópteros de los bosques nativos del Departamento Aluminé (Neuquén-Argentina). — *Bosque, Valdivia* (Chile), **15** (1): 67–74.
- GREHAN, J. R. (1984): The host range of *Aenetus virescens* (Lepidoptera: Hepialidae) and its evolution. — *New Zealand Entomologist*, Nelson, **8**: 52–61.
- (1989): Larval feeding habits of the Hepialidae (Lepidoptera). — *Journal of Natural History*, London, **23**: 803–824.
- (2010): Structural variants in the morphology of the first abdominal tergite supporting the monophyly of the Latin American genera *Cibyra* WALKER, *Druceiella* VIETTE, *Pfitzneriana* VIETTE and *Trichophassus* LE CERF (Lepidoptera: Hepialidae). — *Bulletin of the Buffalo Museum of Science*, Buffalo (N.Y.), **39**: 43–63.
- , & RAWLINS, J. E. (2003): Larval description of a new world ghost moth, *Phassus* sp., and the evolutionary biogeography of wood-boring Hepialidae (Lepidoptera: Exoporia: Hepialoidea). — *Proceedings of the Entomological Society of Washington*, Washington D.C., **105**(3): 733–755.
- HAMPSON, G. F. (1893 [“1892”]): Moths. Vol. 1. — *In*: BLANFORD, W. T. (ed.), *The Fauna of the British India, including Ceylon and Burma*. — London (Taylor & Francis), Calcutta (Thacker, Spink), Bombay (Thacker), Berlin (R. Friedländer), xxxiii + 527 pp. [For the publication year 1893, see OPINION 1698.]
- (1903): Lepidoptera Phalænæ. — *In*: HERON, F. A., & HAMPSON, G. F., *On the Lepidoptera collected at Chapada, Matto Grosso by Mr. A. ROBERT* (Percy Sladen Expedition to Central Brazil). — *Proceedings of the Zoological Society of London*, London, **1903** (2): 259–260.
- HEADLEE, T. J. (1907): A study in butterfly wing-venation, with special regard to the radial vein of the front wing. — *Smithsonian Miscellaneous Collections*, Washington D.C., **48** (3): 284–296, 46 figs., 5 pls.
- HEMMING, F. (1937): HÜBNER. A bibliographical and systematic account of the entomological works of Jacob HÜBNER and of the supplements thereto by Carl GEYER, Gottfried Franz von FRÖLICH and Gottlieb August Wilhelm HERRICH-SCHÄFFER. 2 vols. — London (Royal Entomological Society), xxxiv + 605 + [xi] + 274 pp.
- HERRICH-SCHÄFFER, G. A. W. (1850–[1858]): *Sammlung neuer oder wenig bekannter aussereuropäischer Schmetterlinge*. — Regensburg (G. J. Manz), 84 pp., 120 pls.
- HILJE Q., L., ARAYA F., C., & SCORZA R., F. (1992a): Forest pests in Central America. Field Guide. Technical Manual 4. — Technical series, CATIE (Tropical Agriculture Research and Training Center), Turrialba (Costa Rica), 261 pp.
- , —, —, & VÍQUEZ C., M. (1992b): Forest pests in Central America. Handbook. Technical Manual 3. — Technical series, CATIE (Tropical Agriculture Research and Training Center), English edition, Turrialba (Costa Rica), 187 pp.
- HOFFMANN, F. (1931): Beiträge zur Naturgeschichte brasilianischer Schmetterlinge. II. — *Zeitschrift für wissenschaftliche Insektenbiologie*, Husum, **26** (1): 1–8.
- IHL, E. (1947): *Cuncunilla de los pastos*. — *Simiente*, Temuco (Chile), **17** (1/2): 78–80.
- ISLA M., R. (1959): El control de cuncunilla negra del trébol. — *Agricultura y Ganadería*, Santiago (Chile), **5** (23): 16–18.
- JACKSON S., D. (2007): Observaciones sobre el período de vuelo de *Dalaca chiliensis* (VIETTE, 1950), en la Provincia de Valdivia, Chile (Lepidoptera: Hepialidae). — *Acta Entomológica Chilena*, Santiago de Chile, **31** (1): 35–38.
- JANSE, A. J. T. (1917): Check-list of the South African Lepidoptera Heterocera. — Pretoria (Transvaal Museum), 1–219 + xii pp.
- (1919): Notes on the Hepialid genera *Gorgopis* and *Dalaca*, with descriptions of six apparently new South African species. — *Records of the Albany Museum, Grahamstown*, **3** (3): 233–246.
- (1942): The moths of South Africa 4, Part I. Suborder Jugata. — London (J. Bale & Sons & Staples), Durban (E.P. & Commercial Pr.), xxv + 78 pp, pls. 47–62.
- KIRBY, W. F. (1892): A synonymic catalogue of Lepidoptera Heterocera (moths). Vol. I, Sphinges and Bombyces. — London (Guerney & Jackson), Berlin (R. Friedländer), xii + 951 pp.

- KOÇAK, M. K., & KOÇAK, A. Ö. (2008): First attempt for a list of the patronyms in the LepidopteroLOGY based upon the info-system of the CESA. — Priamus, Ankara, Supplement 12: 1-89.
- KOCH, C. K., & WATERHOUSE, D. F. (eds.) (2000): The distribution and importance of arthropods associated with agriculture and forestry in Chile (Distribucion e importancia de los artropodos asociados a la agricultura y silvicultura en Chile). — ACIAR (Australian Centre for International Agricultural Research), Canberra, Monograph 68: 234 pp.
- KOEBELE, A. (1924): Report on enemies of *Lantana camara* in Mexico, and their introduction into the Hawaiian Islands. — Pp. 54-71 in: PERKINS, R. C. L., & SWEZEY, O. H. (eds.), The introduction into Hawaii of insects that attack *Lantana*. — Bulletin of the Experiment Station of the Hawaiian Sugar Planters' Association, Honolulu, 16: 1-83.
- KRAUSS, N. L. H. (1962): Biological control investigations on *Lantana*. — Proceedings of the Hawaiian Entomological Society, Honolulu, 18 (1): 134-136.
- KRISTENSEN, N. P. (1998): 5. The homoneurous glossata. — Pp. 51-63 in: KRISTENSEN, N. P. (ed.), Lepidoptera, moths and butterflies, 1: Evolution, systematics, and biogeography. — In: FISCHER, M. (serial ed.), Handbook of Zoology 4 (35): Arthropoda: Insecta. — Berlin, New York (W. de Gruyter), x + 491 pp.
- KÜNNETH, F. (1914): Die Stigmenversorgung des Insektenthorax. — Zeitschrift für wissenschaftliche Zoologie, Leipzig, 112 (1): 70-92, 1 pl.
- LE CERF, F. (1919): Observations sur le genre *Phassus* Wlkr.; diagnoses de genres nouveaux et description d'une espèce nouvelle [Lépidopt. Hepialidae]. — Bulletin du Muséum National d'Historie Naturelle, Paris, 25: 469-471.
- LIMA, A. M. COSTA (1936): Terceiro catálogo do insectos que vivem nas plantas do Brasil. — Ministério da Agricultura, Departamento Nacional da Produção vegetal, Escola Nacional de Agronomia, Rio de Janeiro, 460 + iv pp.
- (1945): Insetos do Brasil. Lepidópteros. 5° Tomo, 1ª parte. Série didática no. 7. — Escola Nacional de Agronomia, Rio de Janeiro, 379 pp.
- LLOYD, D. C., & BLACKMAN, R. L. (1966): Investigations of natural enemies of Melolonthinae and Hepialidae. — Report, South American Station, Rio Negro (Argentina, Commonwealth Institute of Biological Control), 20 pp.
- MABILLE, M. P. (1885): Diagnoses de Lépidoptères nouveaux. — Bulletin de la Société Philomathique de Paris, Paris, (7) 9 (2): 55-70.
- (1888): Mission scientifique de Cap Horn. 1882-1883. 6. Zoologie, Insectes, Lépidoptères. — Paris (Gauthier Villars et fils), 35 pp., 3 pls.
- MACÍAS, J., ARGUEDAS, M., HILJE, L., & ZANUNCIO, J. (2005): Plagas forestales neotropicales. — Manejo Integrado de Plagas y Agroecología, Turrialba (Costa Rica), 74 (16): 99-100.
- MALLET, J. (1984): Sex roles in the ghost moth *Hepialus humuli* (L.) and a review of mating in the Hepialidae (Lepidoptera). — Zoological Journal of the Linnean Society, London, 79: 67-82.
- MONTE, O. (1934): Borboletas que vivem em plantas cultivadas. — Boletim de Agricultura, Zootecnia e Veterinária, Série agrícola, Belo Horizonte, 21: iv + 219 + i pp.
- MÖSCHLER, H. B. (1877): Beiträge zur Schmetterlings-Fauna von Surinam. II. — Verhandlungen der kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien, Vienna, 27: 629-700.
- NAIR, K. S. S. (2001): Pest outbreaks in Tropical forest plantations: is there a greater risk for exotic tree species? — Center for International Forestry Research, Desa Putera (Indonesia), vii + 74 pp.
- NÄSSIG, W. A., & SPEIDEL, W. (2007): On the authorships of the Lepidoptera Atlas of the "Reise der Novara", with a list of the taxa of Bombycoidea [s.l.] therein described (Insecta, Lepidoptera, Bombycoidea). — Senckenbergiana biologica, Frankfurt am Main, 87: 63-74.
- NEAVE, A. S. (1939a): A list of the names of genera and subgenera in Zoology from the tenth edition of LINNAEUS 1758 to the end of 1935. — Nomenclator Zoologicus 1, London (Zoological Society of London), xiv + 957 pp.
- (1939b): A list of the names of genera and subgenera in Zoology from the tenth edition of LINNAEUS 1758 to the end of 1935. — Nomenclator Zoologicus 2, London (Zoological Society of London), ii + 1025 pp.
- (1940a): A list of the names of genera and subgenera in Zoology from the tenth edition of LINNAEUS 1758 to the end of 1935. — Nomenclator Zoologicus 3, London (Zoological Society of London), ii + 1065 pp.
- (1940b): A list of the names of genera and subgenera in Zoology from the tenth edition of LINNAEUS 1758 to the end of 1935. — Nomenclator Zoologicus 4, London (Zoological Society of London), ii + 758 pp.
- (1950): Nomenclator Zoologicus 5. — London (Zoological Society of London), viii + 308 pp.
- NIELSEN, E. S., & ROBINSON, G. S. 1983. Ghost moths of southern South America (Lepidoptera: Hepialidae). [With a summary in Spanish prepared by P. GENTILI: »Mariposas Fantasmas del sur de Sudamerica«.] — Entomograph, Copenhagen, 4: 1-192, 431 figs.
- , —, & WAGNER, D. L. (2000): Ghost-moths of the world: a global inventory and bibliography of the Exoporia (Mnesarchaeoidea and Hepialoidea) (Lepidoptera). — Journal of Natural History, London, 34: 823-878.
- , & SCOBLE, M. J. (1986): *Afrotheora*, a new genus of primitive Hepialidae from Africa (Lepidoptera: Hepialoidea). — Entomologica Scandinavica, Copenhagen, 17: 29-54.
- NYE, I. W. B., & FLETCHER, D. S. (1991): Microlepidoptera. — In: NYE, I. W. B. (ed.), The generic names of moths of the world, vol. 6. — London (Trustees of the BMNH), xxix + 368 pp.
- ÖZDIKMEN, H. (2007): New substitute names for three preoccupied lepidopteran genera: *Hercynella* BETHUNE-BAKER, 1893, *Coloneura* DAVIS, 1964 and *Paragorgopis* VIETTE, 1951 (Lepidoptera). — Munis Entomology & Zoology, Ankara, 2 (1): 115-118.
- OTICICA FILHO, J. (1947): Sobre a genitália das fêmeas de Hepialidae (Lepidoptera). — Summa Brasiliensis Biologiae, Rio de Janeiro, 1 (16): 384-428.
- OPINION 1698 (1992): *Brahmaea* WALKER, 1855 (Insecta, Lepidoptera): *Bombyx certhia* FABRICIUS, 1793 confirmed as the type species [Case 2737]. — Bulletin of Zoological Nomenclature, London, 49 (4): 294.
- PACKARD, A. S. (1895): Monograph of the bombycine moths of America north of Mexico, including their transformations and origin of the larval markings and armature. — Memoirs of the National Academy of Sciences, Washington D.C., 7: 5-390, 88 figs., 49 pls., 10 maps.
- PACLT, J. (1944): Zur systematischen Klassifikation der Hepialiden und Nomenklatorisches über dieselbe Familie (Lep.). — Časopis České Společnosti Entomologické, Prague, 41: 141-144.
- (1949): On some Hepialidae (Lepidoptera) from Patagonia. Critical remarks on the reports no. 11 from the LJUNGNER'S Expedition 1932-1934. — Tijdschrift voor Entomologie, s'-Gravenhage, 91: 149-152.

- (1953): Genera of the Hepialidae (Insecta, Lepidoptera). — *Journal of the Asiatic Society, Calcutta*, **19** (2): 141–148.
- (1957): A protest against the atomizing of genera, with special reference to the Hepialidae (Lepidoptera). — *Systematic Zoology, Baltimore*, **6** (1): 51–52.
- PAGENSTECHE, A. (1902): Die arktische Lepidopterenfauna. — Pp. 197–400 in: RÖMER, F., & SCHAUDINN, F. (eds.), *Fauna Arctica. Eine Zusammenstellung der arktischen Tierformen, mit besonderer Berücksichtigung des Spitzbergen-Gebietes auf Grund der Ergebnisse der Deutschen Expedition in das nördliche Eismeer im Jahre 1898. Vol. 2* (2): pp. 1–560, 6 pls. — Jena (G. Fischer).
- PAPE V., H. (2009): *Cuncunilla negra*, alternativas de manejo. — *Cooprinforma, Osorno* (Chile), **93**: 4–6.
- PASTRANA, J. A. (2004): Los Lepidópteros argentinos. Sus plantas hospedadoras y otros sustratos alimenticios. — South American Biological Control Laboratory USDA-ARS; Sociedad Entomológica Argentina, San Miguel de Tucumán (Argentina), viii + 334 pp.
- PFITZNER, R. (1906): Zwei neue Hepialiden aus Süd-Amerika. — *Deutsche Entomologische Zeitschrift "Iris"*, Dresden, **18**: 276–277.
- (1914): Neue Hepialiden. — *Entomologische Rundschau, Stuttgart*, **31** (18): 105–106, (19): 110.
- (1937–1938): Familie: Hepialidae. — Pp. 1289–1302, pls. 99–100 + 185 in: SEITZ, A. (ed.) (1913–1940), *Die Grossschmetterlinge der Erde* **6**. Band. Die Amerikanischen Spinner und Schwärmer. — Stuttgart (A. Kernen), viii pp. + pp. [1–711 + 713–1452 = 2 text parts] + viii pp. + pls. 1–185 [i.e., more than 185 pls.].
- QUAIL, A. (1900): Life histories in the hepialid group of Lepidoptera, with description of one new species, and notes on imaginal structure. — *Transactions of the Entomological Society of London, London*, **1900** (3): 411–432, 2 pls.
- (1903): On the antennae of Hepialidae — Lepidoptera Jugatae. — *Transactions of the Entomological Society of London, London*, **1903** (4): 400–508, 1 pl.
- RAMOS-ELORDUY, J., MORENO, J. M. P., VÁSQUEZ, A. I., LANDERO, I., OLIVA-RIVERA, H., & CAMACHO, V. H. M. (2011): Edible Lepidoptera in Mexico: geographic distribution, ethnicity, economic and nutritional importance for rural people. — *Journal of Ethnobiology and Ethnomedicine, London*, **7** (2): 1–22.
- ROBINSON, G. S. (1977): A taxonomic revision of the genus *Callipielus* BUTLER (Lepidoptera, Hepialidae). — *Bulletin of the British Museum (Natural History), London, Entomology*, **35** (4): 101–121, pls. 1–6.
- , & NIELSEN, E. S. (1984): Hepialidae. — In: HEPPNER, J. B. (ed.), *Atlas of Neotropical Lepidoptera. Checklist: Part 1. Micropterigoidea-Immoidea*. — Den Haag (W. Junk), xxvii + 112 pp.
- RODRÍGUEZ, C., CERDA, L., & PEREDO, H. (1980): Detección de insectos causantes de daños en viveros de *Pinus radiata* de la decima region. — *Bosque, Valdivia*, **3** (2): 73–76.
- ROJAS H., M., & CHACÓN U., P. (1980): *Aepytus (Pseudolaca)* [sic] *serta* (SCHAUS), barrenador del tallo de la curuba. — *Revista Colombiana de Entomología, Bogotá*, **6** (3/4): 63–67.
- SBORDONI, V., & FORESTIERO, S. (1985): *The world of butterflies*. — Dorset, 312 pp., 112 pls.
- SCHAUS, W. (1888): Larva of the large species of Hepialidae, *Phasus triangularis* Hy. EDW., from Vera Cruz, Mex. — *Entomologica Americana, Brooklyn*, **4** (4): 64.
- (1890): New species of Mexican Lepidoptera. — *Entomologica Americana, Brooklyn*, **6** (3): 45–47.
- (1892): Descriptions of new species of Lepidoptera Heterocera from Brazil, Mexico, and Peru. — Part II. — *Proceedings of the Zoological Society of London, London*, **1892**: 318–340.
- (1894): On new species of Heterocera from tropical America. — *Proceedings of the Zoological Society of London, London*, **1894**: 225–243.
- (1901): New species of Heterocera from tropical America — II. — *Journal of the New York Entomological Society, New York*, **9** (2): 73–77.
- (1929): New species of Heterocera (Lepidoptera) from southern Brazil. — *Proceedings of the Entomological Society of Washington, Washington D.C.*, **31** (3): 45–61.
- (1940): 7. New species of British Guiana Heterocera. — *Zoologica, Scientific Contributions of the New York Zoological Society, New York*, **25**: 83–88.
- SCHRÖDER, H. (1967): Die Typen und Typoide des Natur-Museums Senckenberg, **38**: Insecta, Lepidoptera, Hepialidae. — *Senckenbergiana biologica, Frankfurt am Main*, **48** (5/6): 337–344.
- SHEPARD, H. H. (1930): The pleural and sternal sclerites of the Lepidopterous thorax. — *Annals of the Entomological Society of America, Lanham (Md.)*, **23** (2): 237–260, 50 figs.
- SILVA, A. G. A., GONÇALVES, C., GALVÃO, D., GONÇALVES, A., GOMES, J., SILVA, M., & SIMONI, L. (1968): Insetos, hospedeiros e inimigos naturais. Quarto catálogo dos insetos que vivem nas plantas do Brasil. Seus parasitos e predadores. 1° Tomo, parte II. — Rio de Janeiro, GB (Ministério da Agricultura, Departamento de Defesa e Inspeção Agropecuária, Serviço de Defesa Sanitária Vegetal, Laboratório Central de Patologia Vegetal), xxvi + 622 pp.
- SIMONSEN, T. (2002): Wing scale covering supports close relationship between *Callipielus* and *Dalaca*, Austral South American ghost moths (Lepidoptera: Hepialidae). — *Studies on Neotropical Fauna and Environment, Lisse (NL)*, **37** (1): 65–69, 23 figs.
- SNODGRASS, R. E., 1909. The thorax of insects and the articulation of the wings. — *Proceedings of the United States National Museum, Washington D.C.*, **36**: 511–595, 212 figs., 69 pls.
- STAUDINGER, O. (1899): Lepidopteren. — In: *Naturhistorisches Museum, Hamburg* (ed.), *Ergebnisse der Hamburger Magalhaensischen Sammelreise 1892–1893*, **2** (6): 1–117, 1 pl. — Hamburg (L. Friederichsen).
- STRAND, E. (1912a): Exotisch-Lepidopterologisches. — *Archiv für Naturgeschichte, Berlin*, **78A** (9): 143–158.
- (1912b): Sechs neue exotische Grossschmetterlinge. — *Archiv für Naturgeschichte, Berlin*, **77A** (1, Supplement 4): 99–102.
- (1914): Lepidoptera Niepeltiana. Abbildungen und Beschreibungen neuer und wenig bekannter Lepidoptera aus der Sammlung W. NIEPELT. (Ed. W. NIEPELT). — Leipzig (Urban), pp. 1–64 + pls. I–XII (4 col., 8 b.&w.).
- (1916): Lepidoptera Niepeltiana 2. Abbildungen und Beschreibungen neuer und wenig bekannter Lepidoptera aus der Sammlung W. NIEPELT. (Ed. W. NIEPELT). — Zirlau (the editor), pp. [2] + 1–26 + pls. XIII–XVII (b.&w.).
- (1927): Enumération des Hétérocères exotiques (à l'exclusion des Noctuides, des Microlépidoptères, des espèces de l'Île Formosa et des espèces que j'ai décrites dans SEITZ "Macrolépidoptères"). — *Entomologisches Nachrichtenblatt, Troppau*, **1** (3): 42–55.
- SWEZEY, O. H. (1925): Records of introduction of beneficial insects into the Hawaiian Islands. — *Hawaiian Planters' Record, Honolulu*, **29** (4): 369–376.
- TINDALE, N. B. (1941): Revision of the ghost moths (Lepidoptera, Homoneura, family Hepialidae). Part IV. — *Records of the South Australian Museum, Adelaide*, **7**: 15–46, 3 pls.

- (1954): On a new genus of hepialid moths from Rarotonga in the Pacific Islands. — *Annals and Magazine of Natural History*, London, (12) 7: 13–15, 1 pl.
- (1981): The origin of the Lepidoptera relative to Australia. — Pp. 957–976 *in*: KEAST, A. (ed.), *Ecological Biogeography of Australia*. — Den Haag (W. Junk), *Monographiae Biologicae* 41 (2): ix + pp. 609–1434.
- URETA R., E. (1951): Una nueva especie de *Xyleutes* HBN. (Lep. Zeuzeridae). — *Boletín del Museo Nacional de Historia Natural*, Santiago de Chile, 25: 73–77.
- (1956): Nuevos Heteroceros (Lep.) de Chile. — *Boletín del Museo Nacional de Historia Natural*, Santiago de Chile, 26 (6): 271–284.
- (1957): Nuevos Heteroceros (Lep.) de Chile (II parte). — *Boletín del Museo Nacional de Historia Natural*, Santiago de Chile, 27 (2): 154–164, 1 pl.
- VERGARA R., R. A. (2005): IV. Insectos. — *In*: BERNAL E., J. A., & DÍAZ D., C. A. (eds.), *Tecnología para el cultivo de la curuba*. — Manual técnico 6. — Rionegro (Colombia, Corporación Colombiana de Investigación Agropecuaria, CORPOICA; Centro de Investigación La Selva), 180 pp.
- VIETTE, P. (1949a): Contribution à l'étude des Hepialidae (4^e note). Description de deux nouveaux genres sud-américains. — *Revue Française d'Entomologie*, Paris, 16 (1): 52–55.
- (1949b): Contribution à l'étude des Hepialidae (7^e note). Le genre *Trichophassus* LE CERF. — *Bulletin de la Société entomologique de France*, Paris, 54 (5): 72–73.
- (1949c): Contribution à l'étude des Hepialidae (12^e note). Genres et synonymie. — *Lambillionia*, Bruxelles, 49 (9/10): 101–104.
- (1950a): Contribution a l'étude des Hepialidae (11^e note). Sur quelques espèces sud-américaines. — *Annales de la Société entomologique de France*, Paris, 116: 73–81.
- (1950b): Contribution to the study of Hepialidae (9th note): The genus *Phassodes* BETHUNE-BAKER (Lepidoptera). — *Proceedings of the Hawaiian Entomological Society*, Honolulu, 14 (1): 189–190.
- (1950c): Contribution a l'étude des Hepialidae (15^a note). Genres et espèces de l'Amérique latine. — *Revue Française d'Entomologie*, Paris, 17 (1): 52–62.
- (1950d): Contribution a l'étude des Hepialidae (20^e note). Le genre *Druceiella* VRR. — *Bulletin de la Société zoologique de France*, Paris, 75 (4): 165–169.
- (1950e): Contribution a l'étude des Hepialidae (19^{ème} note). — *Entomologisk Tidskrift*, Lund, 71 (2): 144–146.
- (1950f): Contribution à l'étude des Hépiidae (22^{ème} note). Hepialidae du Musée de Leiden. — *Zoologische Mededelingen*, Leiden, 31 (7): 67–77.
- (1951a): Contribution a l'étude des Hepialidae (13^a note) (1). Sur quelques espèces sud-américaines. — *Bulletin Scientifique de Bourgogne*, Dijon, 13: 1–[7].
- (1951b): Contribution à l'étude des Hépiidae (16^e note). Sur quelques espèces de la collection PFITZNER. — *Bulletin de la Société entomologique de France*, Paris, 55 (8): 116–119.
- (1951c): Contribution a l'étude des Hépiidae (Lépid.) (24^{ème} note). — *Bulletin mensuel de la Société Linnéenne de Lyon*, Lyon, 20 (4): 95–96.
- (1951d): Contribution à l'étude des Hepialidae (17^e note). Les genres et leur espèce type. — *Lambillionia*, Bruxelles, 50 (7–10): 73–80.
- (1951e): Contribution to the study of Hepialidae (25th note). On some new or little known species in the British Museum. — *Annals and Magazine of Natural History*, London, (12) 4: 1272–1282.
- (1952a): Contribution à l'étude des Hepialidae (Lepidoptera) (23^{ème} note). — *Annalen des Naturhistorischen Museums Wien*, Vienna, 58: 140–144.
- (1952b): Contribucion al estudio de los Hepialidae. — *Boletín de la Sociedad Venezolana de Ciencias Naturales*, Caracas, 14 (78): 27–30.
- (1953a): Contribution à l'étude des Hepialidae (27^{ème} note). Une nouvelle espèce du Mexique. — *Mitteilungen der Münchner Entomologischen Gesellschaft*, Munich, 42: 20–21.
- (1953b): Contribution to the study of Hepialidae (29th note). — *The Entomologist*, London, 86: 81–82.
- (1956): Contribution à l'étude des Hepialidae (31^{ème} note). Sur quelques spécimens du Tring Museum. — *Transactions of the Royal Entomological Society of London*, London, 107: 373–379.
- (1961): Beiträge zur Kenntnis der Insektenfauna Boliviens XV. Lepidoptera II. Contribution à l'étude des Hepialidae. 34^{ème} note. Espèces sud-américaines nouvelles ou peu connues du Muséum de Munich. — *Opuscula Zoologica*, Munich, 55: 1–7, 6 figs.
- , & FLETCHER, D. S. (1968): The types of Lepidoptera Heterocera described by P. MABILLE. — *Bulletin of the British Museum (Natural History)*, London, Entomology 21 (8): 389–425.
- WAGNER, H., & PFITZNER, R. (1911): Hepialidae. — *In*: AURIVILLIUS, C., & WAGNER, H. (eds.), *Lepidopterorum Catalogus* 4. — Berlin (W. Junk), pp. 1–26.
- WALKER, F. (1856 [10. v.]): List of the specimens of lepidopterous insects in the collection of the British Museum. Part VII. — *Lepidoptera Heterocera*. — London (Trustees of BMNH), [4] pp. + pp. [1509]–1808.
- WALSINGHAM, M. A. (1909–1915): Heterocera. Pyralidina and Hepialina (part.). — *In*: GODMAN, F. C., & SALVIN, O. (eds.), *Biologia Centrali-Americana. Zoology. Insecta. Lepidoptera-Heterocera. Vol. IV. Tineina. Pterophorina. Orneodina, and Pyralidina and Hepialina (part.)*. — London, xii + 482 pp.
- WEYMER, G. (1907): Exotische Lepidopteren. — *Deutsche Entomologische Zeitschrift "Iris"*, Dresden, 20: 1–51.
- , & MAASSEN, J. P. (1890): Lepidopteren gesammelt auf einer Reise durch Columbien, Ecuador, Peru, Brasilien, Argentinien und Bolivien in den Jahren 1868–1877 von Alphons STÜBEL. — Berlin (A. Asher), [1] + xi + 182 pp., 9 pls.
- WILLIAMS, F. X. (1935): *Hepialus pharus* DRUCE. A moth borer attacking sugar cane in Guatemala. — *Hawaiian planters' record*, Honolulu, 39 (4): 292–297.
- WINDER, J. A., & HARLEY, K. L. S. (1983): The phytophagous insects on *Lantana* in Brazil and their potential for biological control in Australia. — *Tropical Pest Management*, London, 29 (4): 346–362.
- ZUKOWSKI, B. (1954): Aegeriidae, Cossidae und Hepialidae (Lep.). — *In*: TITSCHACK, E. (ed.), *Beiträge zur Fauna Perus* 4: 85–94. — Jena (G. Fischer).

Received: 27. ix. 2011, 4. x. 2011